

中国直肠癌手术吻合口漏诊断、预防及处理专家共识(2019版)



扫码阅读电子版

中华医学会外科学分会结直肠外科学组

通信作者:张忠涛, Email: zhangzht@ccmu.edu.cn, 电话: 010-63138475

DOI: 10.3760/cma.j.issn.1671-0274.2019.03.001

吻合口漏(anastomotic leakage, AL)是直肠癌手术常见的严重并发症,其发生率在2.4%~15.9%之间,吻合口漏发生后的病死率可高达16%^[1-3]。吻合口漏不仅影响患者的术后恢复,严重的吻合口漏需再次手术干预,甚至会影响患者的远期生存效果^[4-5]。吻合口漏存在诸多危险因素,例如男性、吸烟、肥胖、肿瘤巨大、患有糖尿病等。近年来,随着全直肠系膜切除(total mesorectal excision, TME)手术的推广、腹会阴联合切除手术数量的减少以及低位(超低位)吻合的增加,加之微创技术的普及、新辅助治疗策略的实施以及器械吻合技术的发展,使得直肠癌术后吻合口漏持续成为结直肠外科的热点问题。为此,中华医学会外科学分会结直肠外科学组发起并组织国内部分结直肠外科专家,以循证为基础、以问题为导向,制定《中国直肠癌手术吻合口漏诊断、预防及处理专家共识(2019版)》,以期国内从事结直肠外科临床工作的同行提供参考。

一、直肠手术吻合口漏的定义、诊断与分级

(一)定义

2010年,直肠手术吻合口漏由国际直肠癌研究组(International Study Group of Rectal Cancer, ISREC)定义为:在结肠-直肠或结肠-肛管吻合部位的肠壁完整性的中断、缺损,使得腔内外间室连通(包括重建直肠储袋缝合线部位的漏,如J-pouch)以及于吻合部位旁出现盆腔脓肿^[6]。(专家赞成率95.45%)

根据吻合口漏发生的时间分为:早期漏(术后30 d内)和迟发漏(术后30 d后)^[7]。(专家赞成率65.91%)

(二)直肠吻合口漏的诊断与分级

吻合口漏分级方法并不统一。2010年,ISREC提出了目前国际较为公认的吻合口漏分级方法,将直肠吻合口漏分为三级,具体见表1^[6]。

表1 国际直肠癌研究组的吻合口漏分级^[6]

分级	临床表现
A级	亚临床吻合口漏,也称作影像学吻合口漏,无临床症状;不需特殊治疗
B级	表现为腹痛、发热,脓性或粪渣样引流物自肛门、引流管或阴道流出(直肠阴道瘘),白细胞及C反应蛋白升高;需保守治疗的吻合口漏
C级	表现为腹膜炎、脓毒症,及其他B级吻合口漏的临床表现;需二次手术治疗的吻合口漏

二、直肠癌手术吻合口漏的相关危险因素

(一)术前因素

1. 性别:男性是术后吻合口漏的独立危险因素^[8-15]。男性患者吻合口漏发生率高于女性患者,与男性骨盆狭窄导致手术难度大、手术时间长相关。(专家赞成率68.18%)

2. 美国麻醉师协会(American Society of Anesthesiologists, ASA)分级:ASA分级与吻合口漏的发生密切相关。ASA分级≥Ⅱ或Ⅲ级被认为是吻合口漏的危险因素^[8-12]。(专家赞成率88.64%)

3. 体质指数(body mass index, BMI):直肠癌术后吻合口漏与高BMI密切相关。BMI≥30 kg/m²显著增加吻合口漏的发生率^[14]。(专家赞成率88.64%)

4. 术前合并症:患者术前有糖尿病、肾功能不全、低蛋白血症等合并症,会增加吻合口漏的发生率^[8,16]。(专家赞成率97.73%)

糖尿病患者的术后吻合口漏发生率明显高于非糖尿病患者,糖尿病为吻合口漏的独立危险因素^[17]。对于肾功能不全的直肠癌患者,特别是在进行急诊手术时,建议谨慎选择一期吻合手术^[16]。术前低白蛋白血症能够反映全身疾病严重程度,可能直接影响吻合口愈合^[18]。(专家赞成率97.73%)

5. 术前肿瘤治疗:对于可能影响吻合口愈合的术前肿瘤治疗,包括新辅助治疗(长程放疗及短程放疗),可能增加直肠癌术后吻合口漏的严重程度,并导致愈合时间的延迟^[19]。(专家赞成率97.73%)

使用抗血管生成的靶向药物(如贝伐单抗)的转化治疗,是否会影响正常组织微循环,从而影响吻合口愈合,仍有争议^[20-21]。(专家赞成率75.00%)

6. 吸烟和饮酒:吸烟和饮酒被认为是吻合口漏的高危因素^[22]。吸烟相关的微血管疾病可能影响结直肠的血供,导致吻合口继发缺血。酗酒可能与营养不良相关,术后易发生心功能不全、免疫抑制及凝血功能不全,是发生吻合口漏的可能机制。(专家赞成率56.82%)

7. 术前药物使用:长期应用糖皮质激素,尤其是联用其他免疫抑制药物,可能增加吻合口漏风险^[23-24]。虽然尚无证据表明,非甾体类抗炎药会增加吻合口漏的发生率,但围手术期应慎重使用该类药物^[25-27]。(专家赞成率100%)

8. 肿瘤状况:肿瘤分期和直径是吻合口漏发生率增加的

危险因素。肿瘤分期和直径的增加,意味着患者的全身状态通常较差,盆腔内手术操作的困难也增加。(专家赞成率56.82%)

研究显示,肿瘤直径 ≥ 5 cm,吻合口漏的发生率增加4倍^[28]。(专家赞成率54.55%)

9.机械性肠道准备和抗生素的使用:机械性肠道准备和抗生素的使用与吻合口漏发生的关系仍有争论^[29-30]。近年来研究认为,机械性肠道准备联合术前口服非肠道吸收性抗生素,可以显著降低吻合口漏的发生率^[31-32]。(专家赞成率56.82%)

(二)术中因素

1.手术方式及入路:直肠癌的手术方式及入路,应根据术者经验、肿瘤位置、分期和患者的身体状况综合决定^[33-35]。众多临床研究及荟萃分析结果显示,不同的吻合方式(端端吻合、端侧吻合等)对于发生吻合口漏的影响无差别(专家赞成率75.00%);腹腔镜手术与开腹手术吻合口漏发生率无显著差异^[36-37]。(专家赞成率86.36%)

TME是中低位直肠癌的金标准术式,由于TME手术完全切除直肠系膜,且在低位离断直肠后实施消化道重建,因此,TME手术比非TME手术的吻合口漏发生率更高^[1]。经肛全直肠系膜切除术(transanal TME, taTME)是近年来新发展起来的术式,欧洲结直肠癌疾病学会(European Society of Coloproctology, ESCP)协作组发布的一项国际多中心临床分析结果显示,taTME术后吻合口漏的发生风险高于腹腔镜辅助TME手术($P=0.02$)^[38]。(专家赞成率54.55%)

2.吻合口与肛缘距离:吻合口与肛缘距离是吻合口漏的重要影响因素,该距离 < 5 cm可将吻合口漏发生风险提高8倍余^[22,39-40]。多因素分析显示,吻合口距肛门距离是吻合口漏发生的独立危险因素^[41]。(专家赞成率93.18%)

3.预防性肠造口:预防性造口对吻合口漏发生率的影响存在争议^[42-46]。预防性造口可以降低吻合口漏所引起的腹膜炎等严重并发症的发生率,也可降低吻合口漏后的再手术率以及吻合口漏相关的病死率。(专家赞成率93.18%)

4.术中出血量与围手术期输血量:术中出血量与吻合口漏的发生具有相关性^[9,28-29]。此外,围手术期输血 ≥ 400 ml被认为是吻合口漏的高危因素^[28]。(专家赞成率52.27%)

5.切断直肠使用闭合器数目:腹腔镜手术中切断直肠使用切割闭合器数目 ≥ 3 个可将吻合口漏的发生风险提高1.42倍;钉合线之间出现缺损增多,导致吻合口漏发生率增加^[28,34,47-49]。(专家赞成率93.18%)

6.其他:保留左结肠动脉,对吻合口漏的发生是否存在影响,仍然存在争议^[28-34,50-51]。(专家赞成率56.82%)

有文献表明,侧方淋巴结清扫会增加术后吻合口漏发生率^[28,52]。(专家赞成率34.09%)

(三)术后因素

术后因素对吻合口漏的影响,主要体现在对患者全身状态调整改善情况方面。即使术前已经纠正贫血、低蛋白血症、高血糖等,经历手术麻醉打击后,术后仍应密切观察患者

的上述各项指标。术后应用非甾体类抗炎药物镇痛是否会增加吻合口漏的发生率,尚缺乏证据。(专家赞成率47.73%)

术后吻合口出血被认为是吻合口漏发生的危险因素,但临床上往往并不能区别是吻合口漏导致了出血,还是吻合口出血继发了吻合口漏。术后早期腹泻可能与直肠癌低位前切除术吻合口漏的发生相关^[53]。(专家赞成率84.09%)

术后关注心肺功能,避免因心肺因素导致的低氧血症也是保证组织灌注,减少吻合口漏发生的必要措施。

三、直肠癌术后吻合口漏的预防

(一)术前预防

积极纠正术前高危因素,对于存在低蛋白水平、糖尿病、贫血、肠梗阻等可能增加吻合口漏风险的患者要在术前积极改善全身状况^[54-56];术前可进行机械性肠道准备联合口服非肠道吸收抗生素^[31-32]。(专家赞成率56.82%)

(二)术中预防

1.预防性造口:预防性造口可以减轻、甚至避免中低位直肠癌前切除术吻合口漏导致的严重腹腔感染、脓肿、感染性休克等危及患者生命的状况发生,避免因吻合口漏导致的二次手术^[57-58]。

中低位直肠癌手术施行预防性造口的手术指征尚有争议,对于如下情况可考虑施行预防性造口术:(1)全身情况较差者;(2)术前存在肠梗阻;(3)存在吻合口漏的高危因素。预防性造口是降低吻合口漏的并发症严重程度、以及再手术风险的方法之一^[45,59]。(依情况做预防性造口,专家赞成率79.55%)

2.吻合口血供:吻合口血供是确保安全吻合的最重要条件之一。为保证吻合口良好的血供,术中应仔细判断边缘动脉避免损伤。在不能确定吻合口血供是否良好的情况下,可以考虑应用术中荧光显影(吲哚菁绿)技术协助判断^[60]。另外,尽管存在争议,诸多研究认为,术中保留左结肠动脉(left colic artery, LCA)可以改善乙状结肠远端的血供^[61-62]。尤其对老年患者,考虑存在动脉硬化等因素时,保留LCA可能是有益的选择。(专家赞成率56.82%)

3.吻合口张力:吻合后的乙状结肠应该是几乎贴附于骶前,避免形成“桥样悬空”,才是吻合口无张力的状态。一般认为,亚洲人的乙状结肠相对较长,手术中不必要常规游离结肠脾曲。但无论是否游离结肠脾曲,都应该保证吻合口彻底无张力。直肠癌根治术中以下情况时应考虑游离结肠脾曲:(1)吻合位置低:超低位吻合、结肠-肛管吻合时,乙状结肠在盆腔内有“桥样悬空”表现;(2)乙状结肠系膜粘连严重,或游离后远端血供欠佳,被迫切除更多肠管;(3)乙状结肠系膜肥厚偏短等;(4)考虑行结肠储袋吻合的情况下。(专家赞成率93.18%)

4.吻合器选择:应根据肠管直径,选择合适的吻合器型号。多数研究认为,圆形吻合器直径与吻合口漏无相关性^[40,49]。横断肿瘤远端直肠的切割闭合器多选择成钉后1.5~1.8 mm的高度,尽量避免使用超过2个切割闭合器。(专家赞成率93.18%)

5. 引流:盆腔引流:多数临床研究或荟萃分析提示,盆腔引流并不能降低吻合口漏的发生^[33,63-64]。但是,盆腔引流可以减少盆腔血肿和感染的发生,减轻吻合口漏的临床症状,并有助于治疗吻合口漏^[39]。一般将引流管放置于吻合口旁以及盆腔的最低处。(常规推荐盆腔引流,专家赞成率 100%)

预置肛管引流:其作用尚存在争议,但诸多研究支持预置肛管在预防直肠癌术后吻合口漏的作用^[65-69];或降低 C 级漏的发生率^[65-67,70-72]。(依情况做预置肛管引流,专家赞成率 52.27%)

6. 吻合质量检测:吻合口检测通常应该采用盆腔注水,经肛门注气的充气试验进行测漏。研究表明,术中肠镜检查有助于判断吻合口质量、吻合口有无出血、吻合口血运情况等。内镜检查并未降低吻合口漏的发生率^[73]。(依情况做吻合口检测,专家赞成率 45.45%)

吻合口缝合加固:吻合口缝合加固能否降低中低位直肠癌术后吻合口漏的发生率存在争议^[74-75]。中低位直肠癌手术多采用双吻合器吻合,吻合口侧方形成两个交角(“狗耳朵区”),此处的缝钉相互交叉,结构薄弱,是吻合口漏的好发部位^[75]。如果操作方便,适当加固缝合是可取的。(依情况做吻合口缝合加固,专家赞成率 54.55%)

四、直肠癌术后吻合口漏的分级处理原则

外科医生应该在遵循基本治疗原则的同时,根据不同等级的吻合口漏,在营养支持治疗及抗感染治疗的基础上,严密观察病情变化,采取个体化治疗策略。对于接受新辅助放疗以及术后需要接受辅助治疗的吻合口漏患者,建议采取更为积极的干预措施。

A 级漏的临床处理对策:无需特殊外科干预,但要在保证引流通畅的前提下,给予全身营养支持治疗以及强有力的抗感染治疗。(专家赞成率 95.45%)

B 级漏的内镜或介入治疗指征:对于吻合口漏较小的患者,可以经过介入途径向盆腔与肛门置管进行双向灌洗、负压吸引以保持吻合口漏周围无粪便聚集。具有通畅引流或吻合口漏出较少的患者,可以试行内镜下治疗,使用覆膜支架对漏口进行封闭^[76-77]。但是,对于距离肛门 <3 cm 的吻合口漏,覆膜支架对肛管刺激强烈常不能耐受;对于吻合口漏直径 >1 cm 者,覆膜支架难以达到促进愈合目的。对考虑愈合时间较长、或者治疗无效的 B 级吻合口漏患者,应积极考虑外科手术干预。(专家赞成率 77.27%)

C 级漏的外科干预:有明显的腹膜炎或出现休克的患者,首选的治疗应该是手术,建议尽早行近端肠管的造口手术或拆除吻合口而实行永久性结肠造口,术中充分冲洗,尽可能清除腹腔内污染物,同时充分引流^[78-80]。(专家赞成率 100%)

编审委员会名单

主任委员:张忠涛

副主任委员:顾晋、王振军、池畔、兰平

成员(按姓名汉语拼音首字母排序):池畔(福建医科大学附属协和医院)、崔滨滨(哈尔滨医科大学附属肿瘤医院)、崔龙

(上海交通大学医学院附属新华医院)、戴勇(山东大学齐鲁医院)、邓海军(南方医科大学南方医院)、丁培荣(中山大学肿瘤防治中心)、董明(中国医科大学第一附属医院)、杜晓辉(解放军总医院)、冯波(上海交通大学医学院附属瑞金医院)、冯勇(中国医科大学附属盛京医院)、傅传刚(同济大学附属东方医院)、高峰(解放军联勤保障部队第九四〇医院)、顾晋(北京大学肿瘤医院、北京大学首钢医院)、韩方海(中山大学孙逸仙纪念医院)、韩加刚(首都医科大学附属北京朝阳医院)、何显力(空军军医大学唐都医院)、胡祥(大连医科大学附属第一医院)、黄忠诚(湖南省人民医院)、姜军(解放军东部战区总医院)、姜可伟(北京大学人民医院)、兰平(中山大学附属第六医院)、李勇(广东省人民医院)、李明(北京大学肿瘤医院)、李心翔(复旦大学附属肿瘤医院)、李永翔(安徽医科大学附属第一医院)、李志霞(首都医科大学附属北京同仁医院)、梁小波(山西省肿瘤医院)、林国乐(中国医学科学院北京协和医学院 北京协和医院)、林建江(浙江大学医学院附属第一医院)、林谋斌(同济大学附属杨浦医院)、刘骞(中国医学科学院肿瘤医院)、刘铜军(吉林大学第二医院)、卢云(青岛大学附属医院)、钱群(武汉大学中南医院)、申占龙(北京大学人民医院)、孙跃明(江苏省人民医院)、所剑(吉林大学第一医院)、童卫东(陆军军医大学大坪医院)、汪欣(北京大学第一医院)、王屹(北京大学人民医院)、王贵英(河北医科大学第四医院)、王磊(中山大学附属第六医院)、王西墨(天津市南开医院)、王锡山(中国医学科学院肿瘤医院)、王振军(首都医科大学附属北京朝阳医院)、王振宁(中国医科大学附属第一医院)、王征(华中科技大学附属湖北协和医院)、王自强(四川大学华西医院)、吴小剑(中山大学附属第六医院)、武爱文(北京大学肿瘤医院)、肖刚(北京医院)、肖毅(中国医学科学院北京协和医学院 北京协和医院)、许剑民(复旦大学附属中山医院)、姚宏伟(首都医科大学附属北京友谊医院)、叶盛威(华中科技大学附属湖北肿瘤医院)、叶颖江(北京大学人民医院)、袁维堂(郑州大学第一附属医院)、张卫(海军军医大学附属长海医院)、钟鸣(上海交通大学附属仁济医院)、周建平(中国医科大学附属第一医院)、周岩冰(青岛大学附属医院)、朱维铭(解放军东部战区总医院)、张忠涛(首都医科大学附属北京友谊医院)

执笔:姚宏伟

参考文献

- [1] Platell C, Barwood N, Dorfmann G, et al. The incidence of anastomotic leaks in patients undergoing colorectal surgery [J]. Colorectal Dis, 2007, 9(1): 71-79.
- [2] Damen N, Spilsbury K, Levitt M, et al. Anastomotic leaks in colorectal surgery [J]. ANZ J Surg, 2014, 84(10): 763-768. DOI: 10.1111/ans.12494.
- [3] Davis B, Rivadeneira DE. Complications of colorectal anastomoses: leaks, strictures, and bleeding [J]. Surg Clin North Am, 2013, 93(1): 61-87. DOI: 10.1016/j.suc.2012.09.014.
- [4] Krarup PM, Nordholm - Carstensen A, Jorgensen LN, et al. Anastomotic leak increases distant recurrence and long-term

- mortality after curative resection for colonic cancer: a nationwide cohort study [J]. *Ann Surg*, 2014, 259 (5): 930-938. DOI: 10.1097/SLA.0b013e3182a6f2fc.
- [5] Sprenger T, Reißbarth T, Sauer R, et al. Long-term prognostic impact of surgical complications in the German Rectal Cancer Trial CAO/ARO/AIO-94 [J]. *Br J Surg*, 2018, 105 (11): 1510-1518. DOI: 10.1002/bjs.10877.
- [6] Rahbari NN, Weitz J, Hohenberger W, et al. Definition and grading of anastomotic leakage following anterior resection of the rectum: a proposal by the International Study Group of Rectal Cancer [J]. *Surgery*, 2010, 147 (3): 339-351. DOI: 10.1016/j.surg.2009.10.012.
- [7] Lim SB, Yu CS, Kim CW, et al. Late anastomotic leakage after low anterior resection in rectal cancer patients: clinical characteristics and predisposing factors [J]. *Colorectal Dis*, 2016, 18 (4): O135-O140. DOI: 10.1111/codi.13300.
- [8] Parthasarathy M, Greensmith M, Bowers D, et al. Risk factors for anastomotic leakage after colorectal resection: a retrospective analysis of 17518 patients [J]. *Colorectal Dis*, 2017, 19 (3): 288-298. DOI: 10.1111/codi.13476.
- [9] 李俊, 安勇博, 吴国聪, 等. 直肠癌前切除术术后吻合口漏的发生率以及影响因素分析 [J]. *中华胃肠外科杂志*, 2018, 21 (4): 413-418. DOI: 10.3760/cma.j.issn.1671-0274.2018.04.011.
- [10] Breugnot AJ, Swets M, Bosset JF, et al. Adjuvant chemotherapy after preoperative (chemo) radiotherapy and surgery for patients with rectal cancer: a systematic review and meta-analysis of individual patient data [J]. *Lancet Oncol*, 2015, 16 (2): 200-207. DOI: 10.1016/S1470-2045 (14) 71199-4.
- [11] Pommergaard HC, Gessler B, Burchardt J, et al. Preoperative risk factors for anastomotic leakage after resection for colorectal cancer: a systematic review and meta-analysis [J]. *Colorectal Dis*, 2014, 16 (9): 662-671. DOI: 10.1111/codi.12618.
- [12] Sciuto A, Merola G, De Palma GD, et al. Predictive factors for anastomotic leakage after laparoscopic colorectal surgery [J]. *World J Gastroenterol*, 2018, 24 (21): 2247-2260. DOI: 10.3748/wjg.v24.i21.2247.
- [13] 陶凯雄, 高金波. 直肠癌术后吻合口漏发生的危险因素 [J]. *中华胃肠外科杂志*, 2018, 21 (4): 384-387. DOI: 10.3760/cma.j.issn.1671-0274.2018.04.005.
- [14] Kawada K, Sakai Y. Preoperative, intraoperative and postoperative risk factors for anastomotic leakage after laparoscopic low anterior resection with double stapling technique anastomosis [J]. *World J Gastroenterol*, 2016, 22 (25): 5718-5727. DOI: 10.3748/wjg.v22.i25.5718.
- [15] Zhou C, Wu XR, Liu XH, et al. Male gender is associated with an increased risk of anastomotic leak in rectal cancer patients after total mesorectal excision [J]. *Gastroenterol Rep (Oxf)*, 2018, 6 (2): 137-143. DOI: 10.1093/gastro/gox039.
- [16] Krysa J, Patel V, Taylor J, et al. Outcome of patients on renal replacement therapy after colorectal surgery [J]. *Dis Colon Rectum*, 2008, 51 (6): 961-965. DOI: 10.1007/s10350-008-9225-4.
- [17] 邓凯, 张建立, 江秀丽, 等. 直肠癌前切除术术后吻合口漏风险因素分析 [J]. *中华胃肠外科杂志*, 2018, 21 (4): 425-430. DOI: 10.3760/cma.j.issn.1671-0274.2018.04.013.
- [18] Kang CY, Halabi WJ, Chaudhry OO, et al. Risk factors for anastomotic leakage after anterior resection for rectal cancer [J]. *JAMA Surg*, 2013, 148 (1): 65-71. DOI: 10.1001/2013.jamasurg.2.
- [19] WAA B, Westerduin E, Aukema TS, et al. Anastomotic leakage and chronic presacral sinus formation after low anterior resection: results from a large cross-sectional study [J]. *Ann Surg*, 2017, 266 (5): 870-877. DOI: 10.1097/SLA.0000000000002429.
- [20] August DA, Serrano D, Poplin E. "Spontaneous," delayed colon and rectal anastomotic complications associated with bevacizumab therapy [J]. *J Surg Oncol*, 2008, 97 (2): 180-185. DOI: 10.1002/jso.20938.
- [21] McDermott FD, Heeney A, Kelly ME, et al. Systematic review of preoperative, intraoperative and postoperative risk factors for colorectal anastomotic leaks [J]. *Br J Surg*, 2015, 102 (5): 462-479. DOI: 10.1002/bjs.9697.
- [22] Kim CW, Baek SJ, Hur H, et al. Anastomotic Leakage after low anterior resection for rectal cancer is different between minimally invasive surgery and open surgery [J]. *Ann Surg*, 2016, 263 (1): 130-137. DOI: 10.1097/SLA.0000000000001157.
- [23] Sliker JC, Komen N, Mannaerts GH, et al. Long-term and perioperative corticosteroids in anastomotic leakage: a prospective study of 259 left-sided colorectal anastomoses [J]. *Arch Surg*, 2012, 147 (5): 447-452. DOI: 10.1001/archsurg.2011.1690.
- [24] Konishi T, Watanabe T, Kishimoto J, et al. Risk factors for anastomotic leakage after surgery for colorectal cancer: results of prospective surveillance [J]. *J Am Coll Surg*, 2006, 202 (3): 439-444. DOI: 10.1016/j.jamcollsurg.2005.10.019.
- [25] Hakkarainen TW, Steele SR, Bastawors A, et al. Nonsteroidal anti-inflammatory drugs and the risk for anastomotic failure: a report from Washington State's Surgical Care and Outcomes Assessment Program (SCOAP) [J]. *JAMA Surg*, 2015, 150 (3): 223-228. DOI: 10.1001/jamasurg.2014.2239.
- [26] Rutegård M, Westermark S, Kverneng HD, et al. Non-steroidal anti-inflammatory drug use and risk of anastomotic leakage after anterior resection: a protocol-based study [J]. *Dig Surg*, 2016, 33 (2): 129-135. DOI: 10.1159/000443216.
- [27] Kverneng Hultberg D, Angenete E, Lydrup ML, et al. Nonsteroidal anti-inflammatory drugs and the risk of anastomotic leakage after anterior resection for rectal cancer [J]. *Eur J Surg Oncol*, 2017, 43 (10): 1908-1914. DOI: 10.1016/j.ejso.2017.06.010.
- [28] Kawada K, Hasegawa S, Hida K, et al. Risk factors for anastomotic leakage after laparoscopic low anterior resection with DST anastomosis [J]. *Surg Endosc*, 2014, 28 (10): 2988-2995. DOI: 10.1007/s00464-014-3564-0.
- [29] Contant CM, Hop WC, van't SHP, et al. Mechanical bowel preparation for elective colorectal surgery: a multicentre randomised trial [J]. *Lancet*, 2007, 370 (9605): 2112-2117. DOI: 10.1016/S0140-6736(07)61905-9.
- [30] Slim K, Vicaute E, Launay-Savary MV, et al. Updated systematic review and meta-analysis of randomized clinical trials on the role of mechanical bowel preparation before colorectal surgery [J]. *Ann Surg*, 2009, 249 (2): 203-209. DOI: 10.1097/SLA.0b013e318193425a.
- [31] Morris MS, Graham LA, Chu DI, et al. Oral antibiotic bowel

- preparation significantly reduces surgical site infection rates and readmission rates in elective colorectal surgery [J]. *Ann Surg*, 2015, 261 (6) : 1034-1040. DOI: 10.1097/SLA.0000000000001125.
- [32] Scarborough JE, Mantyh CR, Sun Z, et al. Combined mechanical and oral antibiotic bowel preparation reduces incisional surgical site infection and anastomotic leak rates after elective colorectal resection: an analysis of colectomy-targeted ACS NSQIP [J]. *Ann Surg*, 2015, 262 (2) : 331-337. DOI: 10.1097/SLA.0000000000001041.
- [33] Nikolian VC, Kamdar NS, Regenbogen SE, et al. Anastomotic leak after colorectal resection: a population-based study of risk factors and hospital variation [J]. *Surgery*, 2017, 161 (6) : 1619-1627. DOI: 10.1016/j.surg.2016.12.033.
- [34] García - Granero E, Navarro F, Cerdán Santacruz C, et al. Individual surgeon is an independent risk factor for leak after double-stapled colorectal anastomosis: an institutional analysis of 800 patients [J]. *Surgery*, 2017, 162 (5) : 1006-1016. DOI: 10.1016/j.surg.2017.05.023.
- [35] Kayano H, Okuda J, Tanaka K, et al. Evaluation of the learning curve in laparoscopic low anterior resection for rectal cancer [J]. *Surg Endosc*, 2011, 25 (9) : 2972-2979. DOI: 10.1007/s00464-011-1655-8.
- [36] Breukink S, Pierie J, Wiggers T. Laparoscopic versus open total mesorectal excision for rectal cancer [J]. *Cochrane Database Syst Rev*, 2006(4) : CD005200.
- [37] Katsuno H, Shiomi A, Ito M, et al. Comparison of symptomatic anastomotic leakage following laparoscopic and open low anterior resection for rectal cancer: a propensity score matching analysis of 1014 consecutive patients [J]. *Surg Endosc*, 2016, 30 (7) : 2848-2856. DOI: 10.1007/s00464-015-4566-2.
- [38] 2017 European Society of Coloproctology (ESCP) collaborating group. An international multicentre prospective audit of elective rectal cancer surgery; operative approach versus outcome, including transanal total mesorectal excision (TaTME) [J]. *Colorectal Dis*, 2018, Suppl 6:S33-S46. DOI:10.1111/codi.14376.
- [39] Qu H, Liu Y, Bi DS. Clinical risk factors for anastomotic leakage after laparoscopic anterior resection for rectal cancer: a systematic review and meta-analysis [J]. *Surg Endosc*, 2015, 29 (12):3608-3617. DOI:10.1007/s00464-015-4117-x.
- [40] Shinji S, Ueda Y, Yamada T, et al. Male sex and history of ischemic heart disease are major risk factors for anastomotic leakage after laparoscopic anterior resection in patients with rectal cancer [J]. *BMC Gastroenterol*, 2018, 18(1) : 117. DOI:10.1186/s12876-018-0846-3.
- [41] Akiyoshi T, Ueno M, Fukunaga Y, et al. Incidence of and risk factors for anastomotic leakage after laparoscopic anterior resection with intracorporeal rectal transection and double-stapling technique anastomosis for rectal cancer [J]. *Am J Surg*, 2011, 202(3) : 259-264. DOI: 10.1016/j.amjsurg.2010.11.014.
- [42] Shiomi A, Ito M, Saito N, et al. Diverting stoma in rectal cancer surgery. A retrospective study of 329 patients from Japanese cancer centers [J]. *Int J Colorectal Dis*, 2011, 26 (1) : 79-87. DOI:10.1007/s00384-010-1036-0.
- [43] Frouws MA, Snijders HS, Malm SH, et al. Clinical relevance of a grading system for anastomotic leakage after low anterior resection: analysis from a national cohort database [J]. *Dis Colon Rectum*, 2017, 60 (7) : 706-713. DOI: 10.1097/DCR.0000000000000800.
- [44] Ilnát P, Guňková P, Peteja M, et al. Diverting ileostomy in laparoscopic rectal cancer surgery: high price of protection [J]. *Surg Endosc*, 2016, 30(11) : 4809-4816. DOI: 10.1007/s00464-016-4811-3.
- [45] Mrak K, Uranitsch S, Pedross F, et al. Diverting ileostomy versus no diversion after low anterior resection for rectal cancer: a prospective, randomized, multicenter trial [J]. *Surgery*, 2016, 159(4) : 1129-1139. DOI:10.1016/j.surg.2015.11.006.
- [46] Wu SW, Ma CC, Yang Y. Role of protective stoma in low anterior resection for rectal cancer: a meta-analysis [J]. *World J Gastroenterol*, 2014, 20(47) : 18031-18037. DOI: 10.3748/wjg.v20.i47.18031.
- [47] Braunschmid T, Hartig N, Baumann L, et al. Influence of multiple stapler firings used for rectal division on colorectal anastomotic leak rate [J]. *Surg Endosc*, 2017, 31(12) : 5318-5326. DOI:10.1007/s00464-017-5611-0.
- [48] Kim JH, Kim HY, Lee IK, et al. Intra-operative double-stapled colorectal or coloanal anastomotic complications of laparoscopic low anterior resection for rectal cancer : double-stapled anastomotic complication could result in persistent anastomotic leakage [J]. *Surg Endosc*, 2015, 29 (11) : 3117-3124. DOI: 10.1007/s00464-014-4035-3.
- [49] Park JS, Choi GS, Kim SH, et al. Multicenter analysis of risk factors for anastomotic leakage after laparoscopic rectal cancer excision: the Korean laparoscopic colorectal surgery study group [J]. *Ann Surg*, 2013, 257 (4) : 665-671. DOI: 10.1097/SLA.0b013e31827b8ed9.
- [50] Boström P, Haapamäki MM, Matthiessen P, et al. High arterial ligation and risk of anastomotic leakage in anterior resection for rectal cancer in patients with increased cardiovascular risk [J]. *Colorectal Dis*, 2015, 17(11) : 1018-1027. DOI: 10.1111/codi.12971.
- [51] Beppu N, Matsubara N, Noda M, et al. A 'high tie' confers an increased risk of anastomotic leakage for lower rectal cancer surgery in patients treated with preoperative radiotherapy [J]. *Surg Today*, 2015, 45(5) : 600-605. DOI: 10.1007/s00595-014-1022-1.
- [52] Koyama M, Murata A, Sakamoto Y, et al. Risk factors for anastomotic leakage after intersphincteric resection without a protective defunctioning stoma for lower rectal cancer [J]. *Ann Surg Oncol*, 2016, 23 Suppl 2:S249-S256. DOI:10.1245/s10434-015-4461-z.
- [53] Hidaka E, Maeda C, Nakahara K, et al. Fecal volume after laparoscopic low anterior resection predicts anastomotic leakage [J]. *Dig Surg*, 2017, 34 (5) : 394-399. DOI: 10.1159/000454960.
- [54] Kingham TP, Pachter HL. Colonic anastomotic leak: risk factors, diagnosis, and treatment [J]. *J Am Coll Surg*, 2009, 208 (2) : 269-278. DOI:10.1016/j.jamcollsurg.2008.10.015.
- [55] Frasson M, Flor-Lorente B, Rodríguez JL, et al. Risk factors for anastomotic leak after colon resection for cancer: multivariate

- analysis and nomogram from a multicentric, prospective, national study with 3193 patients [J]. *Ann Surg*, 2015, 262(2): 321-330. DOI:10.1097/SLA.0000000000000973.
- [56] Schwegler I, von HA, Gutzwiller JP, et al. Nutritional risk is a clinical predictor of postoperative mortality and morbidity in surgery for colorectal cancer [J]. *Br J Surg*, 2010, 97(1):92-97. DOI:10.1002/bjs.6805.
- [57] Shiomi A, Ito M, Maeda K, et al. Effects of a diverting stoma on symptomatic anastomotic leakage after low anterior resection for rectal cancer: a propensity score matching analysis of 1, 014 consecutive patients [J]. *J Am Coll Surg*, 2015, 220(2): 186-194. DOI:10.1016/j.jamcollsurg.2014.10.017.
- [58] Marusch F, Koch A, Schmidt U, et al. Value of a protective stoma in low anterior resections for rectal cancer [J]. *Dis Colon Rectum*, 2002, 45(9): 1164-1171. DOI: 10.1097/01.DCR.0000027040.59190.12.
- [59] Flooden H, Hallböök O, Hagberg LA, et al. Costs and resource use following defunctioning stoma in low anterior resection for cancer - A long-term analysis of a randomized multicenter trial [J]. *Eur J Surg Oncol*, 2017, 43(2): 330-336. DOI:10.1016/j.ejso.2016.12.003.
- [60] Wada T, Kawada K, Takahashi R, et al. ICG fluorescence imaging for quantitative evaluation of colonic perfusion in laparoscopic colorectal surgery [J]. *Surg Endosc*, 2017, 31(10): 4184-4193. DOI: 10.1007/s00464-017-5475-3.
- [61] Seike K, Koda K, Saito N, et al. Laser doppler assessment of the influence of division at the root of the inferior mesenteric artery on anastomotic blood flow in rectosigmoid cancer surgery [J]. *Int J Colorectal Dis*, 2007, 22(6): 689-697. DOI:10.1007/s00384-006-0221-7.
- [62] Fan YC, Ning FL, Zhang CD, et al. Preservation versus non-preservation of left colic artery in sigmoid and rectal cancer surgery: a meta-analysis [J]. *Int J Surg*, 2018, 52:269-277. DOI: 10.1016/j.ijso.2018.02.054.
- [63] Emile SH, Abd ETM. Routine drainage of colorectal anastomoses: an evidence-based review of the current literature [J]. *Gastroenterol Res Pract*, 2017, 2017: 6253898. DOI: 10.1155/2017/6253898.
- [64] Denost Q, Rouanet P, Faucheron JL, et al. To drain or not to drain intraperitoneal anastomosis after rectal excision for cancer: the GRECCAR 5 randomized trial [J]. *Ann Surg*, 2017, 265(3): 474-480. DOI:10.1097/SLA.0000000000001991.
- [65] Matsuda M, Tsuruta M, Hasegawa H, et al. Transanal drainage tube placement to prevent anastomotic leakage following colorectal cancer surgery with double stapling reconstruction [J]. *Surg Today*, 2016, 46(5): 613-620. DOI: 10.1007/s00595-015-1230-3.
- [66] Nishigori H, Ito M, Nishizawa Y. A novel transanal tube designed to prevent anastomotic leakage after rectal cancer surgery: the WING DRAIN [J]. *Surg Today*, 2017, 47(4): 513-520. DOI:10.1007/s00595-016-1392-7.
- [67] Kawada K, Takahashi R, Hida K, et al. Impact of transanal drainage tube on anastomotic leakage after laparoscopic low anterior resection [J]. *Int J Colorectal Dis*, 2018, 33(3): 337-340. DOI:10.1007/s00384-017-2952-z.
- [68] Nishigori H, Ito M, Nishizawa Y, et al. Effectiveness of a transanal tube for the prevention of anastomotic leakage after rectal cancer surgery [J]. *World J Surg*, 2014, 38(7): 1843-1851. DOI:10.1007/s00268-013-2428-4.
- [69] Lee SY, Kim CH, Kim YJ, et al. Impact of anal decompression on anastomotic leakage after low anterior resection for rectal cancer: a propensity score matching analysis [J]. *Langenbecks Arch Surg*, 2015, 400(7): 791-796. DOI:10.1007/s00423-015-1336-5.
- [70] Zhao WT, Hu FL, Li YY, et al. Use of a transanal drainage tube for prevention of anastomotic leakage and bleeding after anterior resection for rectal cancer [J]. *World J Surg*, 2013, 37(1): 227-232. DOI:10.1007/s00268-012-1812-9.
- [71] Yang CS, Choi GS, Park JS, et al. Rectal tube drainage reduces major anastomotic leakage after minimally invasive rectal cancer surgery [J]. *Colorectal Dis*, 2016, 18(12): 0445-0452. DOI: 10.1111/codi.13506.
- [72] Goto S, Hida K, Kawada K, et al. Multicenter analysis of transanal tube placement for prevention of anastomotic leak after low anterior resection [J]. *J Surg Oncol*, 2017, 116(8): 989-995. DOI:10.1002/jso.24760.
- [73] Li VK, Wexner SD, Pulido N, et al. Use of routine intraoperative endoscopy in elective laparoscopic colorectal surgery: can it further avoid anastomotic failure? [J]. *Surg Endosc*, 2009, 23(11): 2459-2465. DOI: 10.1007/s00464-009-0416-4.
- [74] Baek SJ, Kim J, Kwak J, et al. Can trans-anal reinforcing sutures after double stapling in lower anterior resection reduce the need for a temporary diverting ostomy? [J]. *World J Gastroenterol*, 2013, 19(32): 5309-5313. DOI:10.3748/wjg.v19.i32.5309.
- [75] Maeda K, Nagahara H, Shibutani M, et al. Efficacy of intracorporeal reinforcing sutures for anastomotic leakage after laparoscopic surgery for rectal cancer [J]. *Surg Endosc*, 2015, 29(12): 3535-3542. DOI:10.1007/s00464-015-4104-2.
- [76] DiMaio CJ, Dorfman MP, Gardner GJ, et al. Covered esophageal self-expandable metal stents in the nonoperative management of postoperative colorectal anastomotic leaks [J]. *Gastrointest Endosc*, 2012, 76(2): 431-435. DOI:10.1016/j.gie.2012.03.1393.
- [77] Morks AN, Havenga K, Ploeg RJ. Can intraluminal devices prevent or reduce colorectal anastomotic leakage: a review [J]. *World J Gastroenterol*, 2011, 17(40): 4461-4469. DOI:10.3748/wjg.v17.i40.4461.
- [78] 池畔, 陈致奋. 直肠癌术后吻合口漏的诊断与治疗进展 [J]. *中华消化外科杂志*, 2014, 13(7): 584-590. DOI:10.3760/cma.j.issn.1673-9752.2014.07.023.
- [79] Watson AJ, Krukowski ZH, Munro A. Salvage of large bowel anastomotic leaks [J]. *Br J Surg*, 1999, 86(4): 499-500. DOI: 10.1046/J.1365-2168.1999.01096.x.
- [80] Fracalvieri D, Biondo S, Saez J, et al. Management of colorectal anastomotic leakage: differences between salvage and anastomotic takedown [J]. *Am J Surg*, 2012, 204(5): 671-676. DOI:10.1016/j.amjsurg.2010.04.022.

(收稿日期: 2019-01-24)

(本文编辑: 卜建红)