

·诊疗方案·

成人诊断性可弯曲支气管镜检查术
应用指南(2019年版)

中华医学会呼吸病学分会介入呼吸病学学组

通信作者:王广发,北京大学第一医院呼吸和危重症医学科 100034, Email:

wangguangfa@hotmail.com

【摘要】 可弯曲支气管镜检查术是呼吸系统疾病临床诊断和治疗的重要手段,临床应用广泛。本指南是在综合国内外相关文献的基础上,参照国内外相关指南的相关内容,对国内既往指南的内容进行了较多的补充和更新,其目的是规范成人支气管镜检查术的操作,提高疾病的诊断率,降低相关不良风险,减少操作相关的并发症。

DOI: 10.3760/cma.j.issn.1001-0939.2019.08.005

可弯曲支气管镜(包括纤维支气管镜、电子支气管镜,以下简称支气管镜)检查术是呼吸系统疾病临床诊断和治疗的重要手段,临床应用广泛。“成人诊断性可弯曲支气管镜检查术应用指南(2019年版)”(以下简称指南)是在综合国内外相关文献的基础上,按照循证医学的证据等级对相关研究证据进行了分级(表1)。参照国外相关指南的相关内容,对中华医学会呼吸病学分会2008年制定的“诊断性可弯曲支气管镜应用指南(2008年版)”的内容进行了较多的补充和更新。本指南的目的是规范成人支气管镜检查术的操作,提高疾病的诊断率,降低相关不良风险,减少操作相关的并发症。

本指南不涉及以治疗为目的的支气管镜操作。由于硬质气管支气管镜也可用于气管支气管的检查,但其技术迥异于可弯曲支气管镜,且以治疗目的为主,故不在本指南阐述的范畴。

一、支气管镜检查术的适应证及禁忌证

(一)适应证

支气管镜检查术作为临床常用技术,适应证范围非常广泛,对于呼吸系统疾病具有广泛的诊断价值,其中下列情况行支气管镜检查术可显著获益。

1. 疑诊气管、支气管、肺脏肿瘤或肿瘤性病变需要确定病理分型,或确定浸润范围及分期时,应行支气管镜检查术^[2-6](推荐等级B)。鉴于近年来

表1 证据级别及推荐等级^[1]

证据级别	证据类型
1++	随机对照试验高质量的荟萃分析、系统评价,或偏倚可能性很小的随机对照试验
1+	随机对照试验质量较高的荟萃分析、系统评价,或偏倚可能性小的随机对照试验
1-	随机对照试验的荟萃分析、系统评价,或偏倚可能性大的随机对照试验
2++	病例对照或队列研究的高质量系统评价,或出现混杂、偏倚和机遇可能性很小而反映因果关联可能性大的、高质量病例对照或队列研究
2+	出现混杂、偏倚和机遇可能性小而反映因果关联可能性较大的、较高质量的病例对照或队列研究
2-	出现混杂、偏倚和机遇可能性大而反映因果关联可能性明显不足的病例对照或队列研究
3	非分析性研究,即病例报告、系列病例分析
4	专家意见
推荐等级	推荐类型
A	直接适用于目标人群的1++或1+级证据
B	直接适用于目标人群的2++级证据,1++或1+级证据的外推证据
C	直接适用于目标人群的2+级证据,2++级证据的外推证据
D	3或4级证据,2+级证据的外推证据

肺癌靶向治疗、免疫治疗的进展,支气管镜检查术也适用于对肿瘤进行分子病理学诊断和评价,在治疗过程中对病变再活检以对组织病理类型可能的变化及可能继发的基因突变进行评价,以指导后续治疗^[7-10](推荐等级C)。

2. 不明原因咯血持续1周以上的患者,尤其是年龄在40岁以上,即使影像学未见明显异常,仍应行支气管镜检查术以明确出血部位及出血原因^[11-15](推荐等级C)。

3. 对于不能明确诊断、进展迅速、抗菌药物效果欠佳、病变持续存在或吸收缓慢、临床诊断为下呼吸道感染或伴有免疫功能受损的患者,应行支气管镜检查术,并采样行相关病原学检查及某些病原标志物检测,有助于临床的正确诊断或病原学诊断^[16-24](推荐等级C)。

4. 器官或骨髓移植后新发肺部病变,或者疑诊移植抗宿主病、移植肺免疫排斥时,建议行支气管镜检查术协助明确病因^[25-27](推荐等级D)。

5. 临床上难以解释、病情进展或治疗效果欠佳的咳嗽患者,怀疑气管支气管肿瘤、异物或其他病变者,建议行支气管镜检查术^[28-29](推荐等级D)。

6. 原因不明的突发喘鸣、喘息,尤其是固定部位闻及鼾音或哮鸣音,需排除大气道狭窄或梗阻时,建议行支气管镜检查术^[30](推荐等级D)。

7. 对于原因不明的弥漫性肺实质疾病,如间质性肺炎、结节病、肺泡蛋白沉积症及职业性肺病等,均建议行支气管镜检查术进行诊断和鉴别诊断^[31-33](推荐等级D)。

8. 对于可疑气道狭窄的患者,支气管镜检查术是重要的诊断和评价狭窄程度、长度、类型及病因的方法,为进一步治疗提供依据^[34](推荐等级D)。

9. 对于任何原因引起的单侧肺、肺叶或肺段不张,均建议行支气管镜检查术以明确诊断^[35-36](推荐等级D)。

10. 外伤后可疑气道损伤的患者,推荐行支气管镜检查术,以利于明确诊断并评估损伤部位、性质和程度^[37-38](推荐等级D)。

11. 临床症状及影像学表现怀疑各种气管、支气管瘘,如气管食管瘘、支气管胸膜瘘等,均推荐行支气管镜检查术,以确定其病因、部位、大小及类型^[39-41](推荐等级D)。

12. 临床怀疑气道异物者,建议行支气管镜检查术,以确定诊断,评估取出难度,决定治疗方案^[42-44](推荐等级D)。

13. 原因不明的纵隔淋巴结肿大、纵隔肿物等,应行支气管镜检查术,获取病理学标本,进行诊断^[45-47](推荐等级C)。

(二)禁忌证

可弯曲支气管镜检查术应用至今,已积累了丰富的临床经验,目前无绝对禁忌证,其相对禁忌证范围亦日趋缩小。但下列情况行支气管镜检查术时发生并发症的风险显著高于一般人群,检查前应慎重权衡利弊。

1. 急性心肌梗死后4周内不建议行支气管镜检查术;急性心肌梗死后4~6周内若需行支气管镜检查术,建议请心内科医生会诊,充分评估其发生心脏病的风险^[48-50](推荐等级D)。

2. 活动性大咯血时行支气管镜检查术风险较高,若必须行支气管镜检查术时,应做好建立人工气道及急救的准备,以应对出血加重可能导致的窒息^[51-52](推荐等级D)。

3. 血小板计数 $<20\times 10^9/L$ 时不推荐行支气管镜检查术。血小板计数 $<60\times 10^9/L$ 时不推荐行支气管镜下黏膜活检或经支气管肺活检^[53-55](推荐等级D)。

4. 妊娠期间不推荐行支气管镜检查术,若病情需要,除非紧急情况,则尽量推迟至分娩或妊娠28周以后进行,并提前与妇产科医生充分沟通,评估风险^[56](推荐等级D)。

5. 恶性心律失常、不稳定心绞痛、严重心肺功能不全、高血压危象、严重肺动脉高压、颅内高压、急性脑血管事件、主动脉夹层、主动脉瘤、严重精神疾病以及全身极度衰竭等,并发症风险通常较高,若必须行支气管镜检查术时需权衡利弊,应做好抢救准备(推荐等级D)。

二、支气管镜检查术的术前准备及特殊患者的注意事项

(一)患者的告知及知情同意

1. 将支气管镜检查术过程中可能出现的问题向患者提供口头或书面指导,可以提高其对操作的耐受性^[57-62](推荐等级C)。

2. 所有患者在接受检查前需书面告知相关风险,并签署知情同意书^[63-66](推荐等级D)。

(二)术前准备

1. 检查前根据病情,必须拍摄正位X线胸片,或者正侧位X线胸片,或者胸部CT。推荐行胸部CT检查,以便于更精准确定病变部位,有助于决定采样部位及方式^[67-70](推荐等级D)。

2. 若无胃肠动力异常或梗阻,局部麻醉时应在支气管镜检查术前 4 h 开始禁食,术前 2 h 开始禁水;全身麻醉时应在支气管镜检查术前 8 h 开始禁食,术前 2 h 开始禁水^[71-76](推荐等级 B)。

3. 检查前建议建立静脉通道,以方便术中给予镇静及其他药物,并保留至术后恢复期结束^[77](推荐等级 D)。

4. 在检查前不应常规应用抗胆碱能药物(如阿托品等)。该类物质缺乏临床获益证据且存在血流动力学不稳定的潜在风险^[78-82](推荐等级 A)。

5. 对于拟行支气管镜检查术的患者,建议行凝血酶原时间、部分凝血活酶时间、血小板计数检查,以除外严重凝血功能异常^[83-85](推荐等级 D)。

6. 根据“中华人民共和国传染病防治法”、“艾滋病防治条例”及“软式内镜清洗消毒技术规范”等法律法规,检查前应筛查血源性传播疾病,防止医源性感染(推荐等级 D)。

7. 对于有心脏病病史及其危险因素的患者,检查前应行心电图检查^[86-87](推荐等级 D)。

8. 对于拟行活检的患者,推荐提前 5~7 d 停用氯吡格雷,提前 3~5 d 停用替格瑞洛,小剂量阿司匹林可继续使用^[54-55,77-88](推荐等级 C)。

9. 对于需提前停用氯吡格雷或替格瑞洛的患者,若植入冠状动脉药物涂层支架未滿 12 个月或植入冠状动脉金属裸支架未滿 1 个月,则应与心内科医生沟通,共同权衡抗血小板药物使用的利弊;若抗血小板药物治疗方案为氯吡格雷或替格瑞洛联合小剂量阿司匹林,则改为单用小剂量阿司匹林;并于操作第 2 天晨起恢复氯吡格雷或替格瑞洛的使用^[55,89-90](推荐等级 D)。

10. 对于拟行活检的患者,推荐提前 5 d 停用华法林。若术后无明显活动性出血,可在支气管镜检查术后 12~24 h 恢复使用,即操作当天夜里或第 2 天晨起恢复使用^[55,77,89-90](推荐等级 D)。

11. 对于需提前停用华法林的患者,可评估停药期间血栓形成风险(表 2)。若为低风险,则停药期间无需替换为低分子肝素;否则,应替换为低分子肝素抗凝,并于支气管镜操作前 24 h 停药。恢复

华法林使用后仍应继续同时使用低分子肝素直至 INR 达到治疗范围^[55,89-90](推荐等级 D)。

12. 对于拟行活检的患者,达比加群酯及利伐沙班需提前 24 h 停药,不需低分子肝素替换^[89-90](推荐等级 D)。

13. 对疑诊慢性阻塞性肺疾病的患者推荐进行肺功能检查,若通气功能重度减退(FEV_1 占预计值% $<40\%$),建议进行动脉血气分析^[91-92](推荐等级 D)。

14. 慢性阻塞性肺疾病及支气管哮喘患者在支气管镜检查术前应预防性使用支气管舒张剂^[93-97](推荐等级 B)。

15. 吸氧可能升高 $PaCO_2$,因此对于支气管镜检查术前 $PaCO_2$ 已升高者,操作中吸氧可能进一步提高 $PaCO_2$,应警惕,但不需要术前常规进行吸氧试验确定呼吸中枢的敏感性^[98-100](推荐等级 D)。

16. $PaCO_2$ 升高并非静脉应用镇静剂的绝对禁忌证,应充分告知患者及其家属、支气管镜检查医生和麻醉医生存在的潜在风险,应谨慎用药并进行密切监测^[99-101](推荐等级 D)。

17. $PaCO_2$ 升高的患者接受支气管肺泡灌洗术可能导致 $PaCO_2$ 进一步升高,但术后多可自行恢复^[100,102](推荐等级 D)。

(三) 支气管镜检查术的镇静和麻醉

1. 如无禁忌证,应常规给予患者镇静剂^[103-110](推荐等级 B)。

2. 推荐短效苯二氮草类镇静剂咪达唑仑为操作中清醒镇静的首选药物^[74,111-118](推荐等级 C)。

3. 咪达唑仑的具体使用方法:(1)70 岁以下患者的初始剂量推荐为 0.05 mg/kg(不宜超过 3 mg),70 岁以上患者则初始剂量不宜超过 2 mg。在操作开始前 5~10 min 给药,注射后约 2 min 起效;(2)咪达唑仑静脉注射应缓慢,约 1 mg/30 s;(3)如果操作时间长,必要时每次可追加 0.5~1.0 mg,但总量不宜超过 10 mg。年龄 >70 岁、衰弱及慢性病患者应适当减量;(4)本药作用存在较大个体差异,应综合分析患者具体情况,个体化给药^[77,112-113,115-116,119-124](推荐等级 D)。

表 2 血栓形成低风险情况

下肢深静脉血栓或肺栓塞 房颤	下肢深静脉血栓或肺栓塞形成已超过 12 个月,且无易栓症或恶性肿瘤等其他血栓形成的高危因素 CHADS ₂ 评分 ^a 为 0~2 分且无脑卒中或短暂性脑缺血发作病史
心脏机械瓣置换术后	主动脉瓣置换术后,且无房颤及其他脑卒中高危因素(包括糖尿病、高血压、年龄 >75 岁等)

注:^aCHADS₂ 评分为房颤血栓风险评分,其中心功能不全、高血压、年龄 >75 岁、糖尿病各为 1 分,脑卒中为 2 分

4. 丙泊酚镇静效果与咪达唑仑相当,部分研究结果显示患者满意度甚至优于咪达唑仑。但其治疗窗较窄,建议由麻醉科医生或有经验的医生密切进行监测,根据情况随时调整给药速度^[113,115-116,118,125-126](推荐等级C)。

5. 阿片类药物(如芬太尼、舒芬太尼、瑞芬太尼等)可与咪达唑仑、丙泊酚、右美托咪定联合使用,以提高患者对操作的耐受性。操作结束后可根据临床情况积极给予拮抗剂。由于此类药物个体差异大,特别是联合用药时呼吸抑制风险增高,建议由麻醉科医生在监测下给药^[116,118,123,125,127](推荐等级B)。

6. 右美托咪定单药或联合阿片类药物应用可取得良好的镇静效果。多数研究结果提示其镇静效果、患者及支气管镜操作者满意度、脉搏氧饱和度维持等方面均优于咪达唑仑或丙泊酚,但存在苏醒时间延长、血流动力学不稳定等风险。目前推荐由有经验的医生使用;推荐用法为 10~15 min 静脉泵注右美托咪定 0.5~1 $\mu\text{g}/\text{kg}$,维持速度为 0.2~0.7 $\mu\text{g}\cdot\text{kg}^{-1}\cdot\text{h}^{-1}$ ^[128-132](推荐等级C)。

7. 局部麻醉首选利多卡因,且鼻部麻醉推荐使用 2% 利多卡因凝胶^[133-137](推荐等级A)。

8. 行咽喉部麻醉时,推荐使用 1% 利多卡因喷雾,支气管镜通过声带前应局部给药^[77,138-144](推荐等级A)。

9. 行气道麻醉时,首选利多卡因。但因雾化给药气道麻醉效果差,且因药物泄露而导致药物经眼结膜吸收,出现不良反应的比例较高,同时增加利多卡因总用量,故不推荐使用雾化给药方式^[138-141,145](推荐等级C)。

10. 经支气管镜注入利多卡因时,应尽可能减少其用量,以避免心律失常、惊厥等并发症。推荐最大剂量不超过 6~7 mg/kg。对于老年患者、肝功能或心功能损害的患者,使用时应适当减量^[139,146-150](推荐等级C)。

三、支气管镜检查术的术中监护及操作

(一)术中监护及并发症的处理

1. 推荐术中常规监测患者的脉搏氧饱和度^[96-98](推荐等级C)。

2. 术中宜监测患者的心率、心律、呼吸频率及血压^[151-154](推荐等级D)。

3. 有条件时推荐持续监测呼气末二氧化碳分压,其对于呼吸抑制的发现早于脉搏氧饱和度的下降(推荐等级D)。

4. 支气管镜检查室建议配备气管插管及心肺复苏的药品、器械及设备^[77](推荐等级D)。

5. 低氧为支气管镜检查术的常见并发症,但多数呈一过性,通过吸氧易于纠正。推荐术中通过鼻、口或人工气道吸氧。当脉搏氧饱和度明显下降(即 SpO_2 绝对值下降 $>4\%$,或 $\text{SpO}_2 < 90\%$)并持续超过 1 min 时,应积极提高吸氧浓度,必要时停止支气管镜操作,以减少低氧相关损伤的发生^[155-158](推荐等级C)。

6. 支气管镜检查术中,应监测镜下出血情况,可根据表 3 判断出血程度,并给予相应处理^[77](推荐等级D)。

表 3 支气管镜操作中出血程度分级及相应处理方式

出血程度	相应处理
无出血	无需持续吸引,出血可自发停止
轻度出血	需持续吸引,出血可自发停止
中度出血	需以支气管镜阻塞活检的叶段支气管,局部使用肾上腺素或冰盐水止血
重度出血	需放置支气管阻塞球囊或导管、外科介入,使用全身凝血剂
极重度出血	可导致输血、窒息、插管、心肺复苏或者死亡,需进入重症监护室

7. 支气管镜检查术后气胸的总体发生率约为 0.1%。但经支气管肺活检(transbronchial lung biopsy, TBLB)后气胸发生率可达 1%~6%,但 TBLB 术后无需常规行胸片检查。若患者出现相关症状,临床怀疑气胸时则应尽快拍摄胸片以确定或排除诊断^[159-163](推荐等级C)。

8. 支气管镜检查术前预防性使用抗菌药物并无获益,即使对有脾切除、感染性心内膜炎病史患者等特殊情况也不例外^[164-170](推荐等级C)。

9. 支气管镜检查术所致菌血症的发生率约为 6%。术后部分患者可因肺泡巨噬细胞释放的某些炎性介质出现一过性发热,其发生率约为 5%~10%,通常不需要进行特殊处理,但应与术后感染进行鉴别^[164,166-167,169](推荐等级D)。

(二)诊断性支气管镜检查术操作的实施标准

1. 对于镜下所见新生物活检时,如无特殊情况,5 块活检标本可满足病理免疫组织化学染色及基因检测需要,保证诊断率^[171-176](推荐等级B)。

2. 对于镜下所见支气管黏膜呈浸润性病变或高度怀疑肿瘤时,推荐联合进行活检、刷检和冲洗,且应在其他操作后进行冲洗,以提高阳性率^[171,173,177-182](推荐等级B)。

3. 高度怀疑肿瘤或癌前病变时,有条件者可考虑在普通光支气管镜检查的基础上结合荧光支气管镜或窄谱成像支气管镜检查,以提高发现病变的敏感度^[171,183-184](推荐等级D)。

4. 对于支气管腔外病变,推荐经支气管针吸活检(transbronchial needle aspiration, TBNA)或支气管腔内超声引导下的经支气管针吸活检(endobronchial ultrasound-transbronchial needle aspiration, EBUS-TBNA)以提高阳性率。传统TBNA操作时可进行快速现场评价(rapid on-site evaluation, ROSE)以减少穿刺针数、评估样本中肿瘤细胞数量及质量。研究表明,ROSE可降低传统TBNA的并发症,但并不影响诊断率^[45-47,185-193](推荐等级C)。

5. 周围型肺部病变患者行活检时,建议应用X线透视、电磁导航、虚拟导航、径向支气管内超声、超细支气管镜等手段,以提高诊断阳性率^[171,194-200](推荐等级C)。

6. 弥漫性肺部病变患者行活检时,无需常规应用X线透视^[201-206](推荐等级B)。

7. 对弥漫性肺部病变患者进行TBLB时,推荐尽可能从一侧肺取4~6块标本,不推荐同时进行双侧肺活检,以避免双侧同时出现严重并发症,导致治疗困难,或无法判断严重并发症的部位而影响紧急处置^[201-203,207](推荐等级C)。

8. 对弥漫性肺部病变或外周型病变患者,经支气管冷冻肺活检可提供更大、质量更高的组织样本,特别是避免了TBLB时对组织的挤压,造成病理判读上的困难。但本操作可能增加气胸及严重出血的风险,推荐在全身麻醉或深度镇静下通过硬质气管支气管镜或气管插管进行^[208-213](推荐等级D)。

9. 对疑诊结节病的患者,推荐进行黏膜活检、TBLB联合支气管肺泡灌洗液的CD₄⁺/CD₈⁺比例检测。若纵隔淋巴结增大,还可考虑联合TBNA或EBUS-TBNA以增加诊断阳性率^[214-221](推荐等级C)。

10. 对免疫功能受损的患者,若存在肺部浸润影,推荐常规行支气管镜检查术,进行刷检、支气管肺泡灌洗术及TBLB,获取标本进行病原学检测,特别是分枝杆菌、真菌(包括肺孢子菌)和病毒(尤其是巨细胞病毒)检测^[222-237](推荐等级C)。

11. 我国为结核病高流行地区,支气管镜检查术获取的标本推荐常规进行抗酸杆菌检测;高度怀疑结核分枝杆菌感染的患者推荐常规进行结核分

枝杆菌培养,并于支气管镜检查术后常规进行痰标本的相关检查^[238-247](推荐等级C)。

12. 对疑诊侵袭性肺曲霉病的患者,应进行支气管肺泡灌洗液镜检及真菌培养;应进行支气管肺泡灌洗液半乳甘露聚糖(galactomannan, GM)测定,该项检查对肺曲霉病的诊断具有较高的敏感度和特异度;由于活检出血风险较高,应根据临床情况权衡利弊,确定是否行TBLB和(或)黏膜活检^[248-256](推荐等级C)。

13. 对于疑诊社区获得性肺炎的患者,疗效不佳或病情迅速进展时,建议在条件许可的情况下进行支气管肺泡灌洗液嗜肺军团杆菌聚合酶链式反应(polymerase chain reaction, PCR)和其他常见病原体的相关检测^[257-265](推荐等级D)。

四、在重症监护室实行的支气管镜检查术

1. 重症监护室患者行支气管镜检查术并发症的发生率高于一般患者^[266-268](推荐等级D)。

2. 支气管镜检查过程中及检查后,应对患者进行连续的多导生命体征监测^[269-275](推荐等级D)。

3. 对于需呼吸机(包括无创呼吸机及有创呼吸机)辅助通气的患者应采取积极措施,如提高吸入氧浓度,将支气管镜通过三通接口插入气管导管内,保证支气管镜检查过程中维持足够的通气和氧合^[269,271,276-279](推荐等级D)。

4. 有以下情况的患者进行操作的风险较高,检查前需谨慎权衡利弊:(1)机械通气时呼气末正压(positive end expiratory pressure, PEEP)>14 cmH₂O(1 cmH₂O=0.098 kPa)、不能耐受分钟通气量减少或检查前依赖高浓度氧疗;(2)颅内高压;(3)气管插管的内径与支气管镜外径差值<2 mm^[266-268,271,279-285](推荐等级D)。

5. 对于肺叶切除术后的机械通气患者,强烈不推荐常规进行支气管镜检查术及支气管肺泡灌洗术来预防肺不张^[286-290](推荐等级A)。

6. 疑诊呼吸机相关性肺炎的患者,强烈建议优先使用非侵入性检查手段以获得病原学证据,仅上述方法无效时,才考虑行支气管镜检查术^[291-306](推荐等级A)。

7. 经可弯曲支气管镜引导下气管插管,可在镜下观察并引导气管插管至恰当位置,同时观察有无气管损伤、出血、感染以及分泌物的情况。对于有颈椎损伤的患者,可弯曲支气管镜引导插管可在颈椎自然位置下进行,避免头颈部伸屈活动。对于颈椎有不稳定骨折、脱位的患者,可避免因气管插管

导致颈椎进一步损伤^[307-309](推荐等级 D)。

8. 呼吸机辅助通气患者进行 TBLB 操作时容易出现气胸、出血、一过性血压下降等并发症,故 TBLB 操作前应充分评估临床获益及风险。但其总体并发症发生率和操作相关的病死率并未显著升高^[310-312](推荐等级 D)。

五、术后处理

1. 局部麻醉结束 2 h 后或全身麻醉结束 6 h 后方可进食、饮水,以避免因咽喉仍处于麻醉状态而导致误吸^[72-74,76](推荐等级 D)。

2. 应通过口头或书面形式告知已行 TBLB 的患者,离院后仍可能发生气胸,如出现憋气、胸疼等症状时应及时就诊^[161-163](推荐等级 D)。

3. 对使用镇静剂的患者,应口头或书面告知其在 24 h 内不要驾车、签署法律文件或操作机械设备^[313-314](推荐等级 D)。

4. 使用镇静剂的门诊患者,应有人陪伴回家,避免自行驾车。对于老年人或行 TBLB 的高危患者,当日应有人在家中陪同^[77,315](推荐等级 D)。

5. 支气管镜检查术后,若为局部麻醉下操作推荐至少观察 30 min;若为全身麻醉,推荐至少观察 6 h,并判断患者生命体征平稳,无意识异常、呼吸困难、胸痛及咯血等情况,方可离院^[316-317](推荐等级 D)。

六、支气管镜的清洗、消毒及医务人员的防护

支气管镜的清洗、消毒以及医务人员的防护应参照中华人民共和国国家卫生健康委员会最新发布的“软式内镜清洗消毒技术规范”^[318],综合近年来相关指南及规范^[77,319-327],提出如下建议。

(一)支气管镜的清洗和消毒

1. 检查开始前、所有检查完成后及 2 名受检者检查之间,均应对支气管镜进行清洗和消毒(推荐等级 D)。

2. 支气管镜的清洗和消毒应由经过培训的专业人员在内镜洗消间内进行(推荐等级 D)。

3. 应每日监测使用中的消毒剂的有效浓度,并保存记录,低于有效浓度时应立即更换(推荐等级 D)。

4. 清洁过程的第一步亦最重要的一步:应用医用清洗剂彻底清洗支气管镜(推荐等级 B)。

5. 每次使用后应更换清洗剂,清洁毛刷宜使用一次性产品,重复使用的毛刷在使用后应进行灭菌或高度消毒水平的清洁处理(推荐等级 C)。

6. 戊二醛对分枝杆菌杀灭作用起效较慢,过氧

乙酸、二氧化氯和过氧化氢则起效较快(≤ 5 min),且较戊二醛的刺激性小,但较容易损伤支气管镜和清洗器具,稳定性较差,且价格较贵(推荐等级 C)。

7. 采用 2% 戊二醛进行手工或自动消毒时,支气管镜的浸泡时间不得少于 20 min(推荐等级 C)。

8. 确诊或疑诊分枝杆菌感染患者使用过的内镜及其附件,其消毒时间应遵循消毒产品的使用说明。确诊或疑似人类免疫缺陷病毒(human immunodeficiency virus, HIV)感染患者使用过的内镜及附件,目前暂无特殊消毒规定。

9. 分枝杆菌(如龟分枝杆菌)抵抗力强,建议使用含氯消毒剂或过氧乙酸消毒剂(推荐等级 B)。

10. 对乙型肝炎、HIV 阳性以及怀疑结核病的患者,应安排在最后进行检查(推荐等级 D)。

11. 为最大限度减少工作人员与消毒剂及消毒剂挥发气体的接触,推荐使用自动清洗消毒机对支气管镜进行清洗和消毒(推荐等级 C)。

12. 自动清洗消毒机必须设有消毒槽、浸洗盘和各种液体通道,并常规对自动洗镜机及其配件进行检测和消毒(推荐等级 C)。

13. 推荐使用灭菌水或过滤水对消毒后的内镜进行最终漂洗(推荐等级 D)。

14. 若冲洗用水的质量难以保证时,应采用 75%~95% 的乙醇擦洗支气管镜的外表面,并冲洗管腔,这样可以杀灭包括分枝杆菌在内的非芽孢菌,且乙醇挥发后管腔会迅速干燥。在每次检查完毕及支气管镜存放前,推荐使用这种方法(推荐等级 C)。

15. 对于热稳定的部件或配件(如活检钳)需用机械清洗装置(如超声清洗机)进行清洗,然后进行高压灭菌或其他灭菌处理(推荐等级 D)。

16. 所有冲洗用水通道(水槽、过滤器及管道)的设计都应方便常规清洗和消毒(推荐等级 D)。

17. 支气管镜室应建立内镜清洗消毒登记制度,登记内容应包括患者姓名、使用内镜及其他重复使用的器械编号、清洗时间、消毒时间以及操作人员姓名等事项,宜开展清洗消毒的信息化管理,做到可追溯(推荐等级 D)。

18. 储存支气管镜前需在专门的地点进行干燥,最好装备空气干燥设备。(推荐等级 D)。

19. 支气管镜必须悬挂储存,并保持环境干燥(推荐等级 D)。

20. 建立工作人员培训制度,当引进新型号的支气管镜或处理设备时,一定要准备相应的技术说

明书,并进行培训(推荐等级D)。

(二)医务人员的防护

1. 当怀疑有污染时,培养范围必须包括支气管镜及其器械、自来水及清洗、消毒处理设备(推荐等级C)。

2. 当怀疑有感染发生时,应向医院感染管理部门、支气管镜生产商、疾病预防和控制中心及卫生行政部门通报情况(推荐等级C)。

3. 所有医务人员应接种乙型肝炎疫苗,在适当的时候检测机体的免疫状态(推荐等级D)。

4. 在行支气管镜检查术过程中,医务人员应穿戴防护用具,包括隔离衣或防水围裙、口罩、护目镜和手套(推荐等级C)。

5. 对确诊或疑诊多重耐药结核分枝杆菌感染的患者进行支气管镜检查术时,医务人员推荐佩戴医用防护口罩(推荐等级D)。

6. 医务人员所使用的手套应不含滑石粉(推荐等级B)。

7. 针状活检钳等锐利附件的清洗应格外小心,以防止医务人员刺伤(推荐等级C)。

8. 工作中可能与醛类物质接触的所有医务人员均应在参加工作前进行体检;参加工作后,职业保健部门应定期检查其肺功能,了解其有无不适主诉(推荐等级D)。

9. 为了尽可能避免医务人员与消毒剂接触,支气管镜最好在装有自动通风系统的专用房间内消毒,有条件者在烟尘柜中进行更好(推荐等级D)。

10. 在清洗和消毒器械过程中,医务人员应穿戴防护用具,包括丁腈橡胶手套、能保护双眼的护目镜或防护面屏、口罩以及塑料隔离衣,以免受到溅出的污水、雾化液和蒸汽的侵害(推荐等级D)。

11. 使用一次性附件(尤其是注射针)可以减少医务人员在清洁器械过程中被感染的风险(推荐等级C)。

12. 为了避免医务人员与消毒剂接触,应尽可能使用高压蒸汽灭菌器械或一次性器械(推荐等级C)。

13. 从事支气管镜操作的专业人员,应接受有关患者护理、感染控制、器械清洁(包括醛类物质的安全使用及在器械清洁过程中可能危害健康的因素)等知识的培训(推荐等级D)。

14. 经常使用透视或CT辅助进行支气管镜检查术操作的人员要求其左侧胸上部常年佩戴放射剂量检测仪,保持清洁,防止污染,时间不超过

90 d(推荐等级D)。

15. 对职业照射人员个人规定的剂量限值:(1)成年人连续5年的年平均有效剂量为20 mSv,但不可做任何追溯性年平均;连续5年中的任何单一年份的年平均有效剂量为50 mSv,但连续5年平均有效剂量不得超过20 mSv;眼部晶状体的年当量剂量为150 mSv;四肢或皮肤的年当量剂量为500 mSv。(2)确认怀孕后,职业照射人员将执行与公众相同剂量限值。(3)在特殊情况下,可以对个人年剂量限值做下述临时改变:按审管部门规定,连续5年的平均期可破例延长至10个连续年;10年内任何一位职业照射人员个人的年平均有效剂量不得超过20 mSv;在10个连续年期间任何单一年份受到的年平均有效剂量不得超过50 mSv;在10个连续年期间,自延长期以来任何一位职业照射人员受到的有效剂量累计达到100 mSv时,应对此进行审查。对个人剂量限值的临时变更应遵守审管部门的规定,任何一年内不得超过50 mSv;临时的改变期限不得超过5年(推荐等级D)。

16. 操作环境需要符合国家放射防护标准的要求,有安全设置、电离辐射警告标志、照射状态指示灯、“门-机”连锁装置和防辐射措施(推荐等级D)。

顾问:陈荣昌、瞿介明

执笔人:黄璐君、章巍、王广发

编写组成员(排名不分先后):王广发、李时悦、李强、张杰、曾奕明、柯会星、王洪武、王臻、杨震、赵静、蔡志刚、侯刚、尹金植、白冲、任涛、孙加源、王昌惠、项轶、张新、蒋军红、陈成水、陈恩国、马文江、叶乐平、柯明耀、许飞、林殿杰、马芸、杨华平、周锐、陈正贤、冯起校、李静、荣福、柳广南、郭述良、王可、叶贤伟、邢西迁、金发光、周玮、程渊、张红、于鲲鹏、余勃、胡展维、刘亚男、黄俊芳、郭翠艳、唐海燕、王东信、李六亿

利益冲突 所有作者均声明不存在利益冲突

参 考 文 献

- [1] Harbour R, Miller J. A new system for grading recommendations in evidence based guidelines[J]. BMJ, 2001, 323(7308): 334-336. DOI: <http://dx.doi.org/10.1136/bmj.323.7308.334>.
- [2] Mazzone P, Jain P, Arroliga AC, et al. Bronchoscopy and needle biopsy techniques for diagnosis and staging of lung cancer[J]. Clin Chest Med, 2002, 23(1): 137-158, ix. DOI: 10.1016/S0272-5231(03)00065-0.
- [3] Arroliga AC, Matthay RA. The role of bronchoscopy in lung cancer[J]. Clin Chest Med, 1993, 14(1): 87-98.
- [4] Hyer JD, Silvestri G. Diagnosis and staging of lung cancer[J]. Clin Chest Med, 2000, 21(1): 95-106, viii-ix. DOI: 10.1016/

- S0272-5231(05)70010-1.
- [5] Rafanan AL, Mehta AC. Role of bronchoscopy in lung cancer [J]. *Semin Respir Crit Care Med*, 2000, 21(5): 405-420. DOI: 10.1055/s-2000-9408.
 - [6] Wang KP. Staging of bronchogenic carcinoma by bronchoscopy [J]. *Chest*, 1994, 106(2): 588-593. DOI: 10.1378 / chest.106.2.588.
 - [7] Kang YR, Park HY, Jeon K, et al. EGFR and KRAS mutation analyses from specimens obtained by bronchoscopy and EBUS-TBNA[J]. *Thorac Cancer*, 2013, 4(3): 264-272. DOI: 10.1111/1759-7714.12006.
 - [8] Castro RL, Borrego H, Disdier C, et al. Feasibility of bronchoscopy obtained tissue for tumor diagnosis and testing epidermal growth factor receptor (EGFR) mutational status[J]. *European Respiratory Journal*, 2013, 42(Suppl 57):4532.
 - [9] Yamamoto G, Kikuchi M, Kobayashi S, et al. Routine genetic testing of lung cancer specimens derived from surgery, bronchoscopy and fluid aspiration by next generation sequencing[J]. *Int J Oncol*, 2017, 50(5): 1579-1589. DOI: 10.3892/ijo.2017.3935.
 - [10] Tsai TH, Yang CY, Ho CC, et al. Multi-gene analyses from waste brushing specimens for patients with peripheral lung cancer receiving EBUS-assisted bronchoscopy [J]. *Lung Cancer*, 2013, 82(3): 420-425. DOI: 10.1016 / j. lungcan.2013.10.005.
 - [11] Hirshberg B, Biran I, Glazer M, et al. Hemoptysis: etiology, evaluation, and outcome in a tertiary referral hospital [J]. *Chest*, 1997, 112: 440-444. DOI: 10.1378/chest.112.2.440.
 - [12] Jackson CV, Savage PJ, Quinn DL. Role of fiberoptic bronchoscopy in patients with hemoptysis and a normal chest roentgenogram[J]. *Chest*, 1985, 87(2):142-144. DOI: 10.1378/ chest.87.2.142.
 - [13] Poe RH, Israel RH, Marin MG, et al. Utility of fiberoptic bronchoscopy in patients with hemoptysis and a nonlocalizing chest roentgenogram [J]. *Chest*, 1988, 93(1): 70-75. DOI: 10.1378/chest.93.1.70.
 - [14] Gong H Jr, Salvatierra C. Clinical efficacy of early and delayed fiberoptic bronchoscopy in patients with hemoptysis [J]. *Am Rev Respir Dis*, 1981, 124(3): 221-225. DOI: 10.1164/ arrd.1982.125.2.269a.
 - [15] McGuinness G, Beacher JR, Harkin TJ, et al. Hemoptysis: prospective high-resolution CT/bronchoscopic correlation [J]. *Chest*, 1994, 105(4): 1155-1162. DOI: 10.1378 / chest.105.4.1155.
 - [16] Helmers RA, Pisani RJ. Bronchoalveolar lavage. In: Prakash UB, ed. *Bronchoscopy*[J]. New York, NY: Raven Press, 1994: 155-182.
 - [17] Reynolds HY. Bronchoalveolar lavage [J]. *Am Rev Respir Dis*, 1987, 135: 250-263. DOI:10.1164/arrd.1987.135.1.250.
 - [18] Guidelines for the management of adults with hospital-acquired, ventilator-associated, and health-care-associated pneumonia [J]. *Am J Respir Crit Care Med*, 2005, 171(4): 388-416. DOI: 10.1164 / rccm.200405-644ST.
 - [19] Campbell JH, Blessing N, Burnett AK, et al. Investigation and management of pulmonary infiltrates following bone marrow transplantation: an eight year review [J]. *Thorax*, 1993, 48(12): 1248-1251. DOI:10.1016/0036-9748(84)90327-2.
 - [20] Rano A, Agusti C, Jimenez P, et al. Pulmonary infiltrates in non-HIV immunocompromised patients: a diagnostic approach using non-invasive and bronchoscopic procedures [J]. *Thorax*, 2001, 56(5): 379-387. DOI:10.1136/thorax.56.5.379.
 - [21] Hobenadel IA, Kiworr M, Genitsariotis R, et al. Role of bronchoalveolar lavage in immunocompromised patients with pneumonia treated with a broad spectrum antibiotic and antifungal regimen [J]. *Thorax*, 2001, 56(2): 115-120. DOI: 10.1136/thorax.56.2.115.
 - [22] Chastre J, Wolff M, Fagon JY, et al. Comparison of 8 vs 15 days of antibiotic therapy for ventilator-associated pneumonia in adults: a randomized trial[J]. *Jama*, 2003, 290(19): 2588-2598. DOI:10.1001/jama.290.19.2588.
 - [23] Jain S, Self WH, Wunderink RG, et al. Community-acquired pneumonia requiring hospitalization among US adults[J]. *New England Journal of Medicine*, 2015, 373(5): 415-427. DOI: 10.1056/NEJMoa1405870.
 - [24] Kalil AC, Metersky ML, Klompas M, et al. Management of adults with hospital-acquired and ventilator-associated pneumonia: 2016 clinical practice guidelines by the Infectious Diseases Society of America and the American Thoracic Society[J]. *Clinical Infectious Diseases*, 2016, 63(5): e61-e11. DOI:10.1093/cid/ciw353.
 - [25] McWilliams TJ, Williams TJ, Whitford HM, et al. Surveillance bronchoscopy in lung transplant recipients: risk versus benefit [J]. *Heart Lung Transplant*, 2008, 27(11): 1203-1209. DOI: 10.1016/j.healun.2008.08.004.
 - [26] Smith L, Singer JP, Hayes M, et al. An analysis of potential risk factors for early complications from fiberoptic bronchoscopy in lung transplant recipients[J]. *Transpl Int*, 2012, 25(2): 172-178. DOI: 10.1111/j.1432-2277.2011.01392. x.
 - [27] Hopkins PM, Aboyoun CL, Chhajed PN, et al. Prospective analysis of 1, 235 transbronchial lung biopsies in lung transplant recipients[J]. *J Heart Lung Transplant*, 2002, 21 (10): 1062-1067. DOI: 10.1016/s1053-2498(02)00442-4.
 - [28] Palombini BC, Villanova CA, Araújo E, et al. A pathogenic triad in chronic cough: asthma, postnasal drip syndrome, and gastroesophageal reflux disease[J]. *Chest*, 1999, 116(2): 279-284. DOI: 10.1378/chest.116.2.279.
 - [29] Irwin RS, Curley FJ, French CL. Chronic cough. The spectrum and frequency of causes, key components of the diagnostic evaluation, and outcome of specific therapy[J]. *The American review of respiratory disease*, 1990, 141(3): 640-647.
 - [30] Mise K, Savicevic AJ, Pavlov N, et al. Removal of tracheobronchial foreign bodies in adults using flexible bronchoscopy: experience 1995-2006[J]. *Surgical endoscopy*, 2009, 23(6): 1360-1364. DOI: 10.1007/s00464-008-0181-9.
 - [31] Meyer KC, Raghu G, Baughman RP, et al. An Official American Thoracic Society Clinical Practice Guideline: The Clinical Utility of Bronchoalveolar Lavage Cellular Analysis in Interstitial Lung Disease[J]. *Am J Respir Crit Care Med*, 2012, 185(9): 1004-1014. DOI: 10.1164/rccm.201202-0320st.
 - [32] Hagmeyer L, Theegarten D, Wohlschläger J, et al. The role of transbronchial cryobiopsy and surgical lung biopsy in the diagnostic algorithm of interstitial lung disease[J]. *The clinical respiratory journal*, 2016, 10(5): 589. DOI: 10.1111/crj.12261.
 - [33] Morales MM, Souza SAL, Loivos LP, et al. Pilot safety study of intrabronchial instillation of bone marrow-derived mononuclear cells in patients with silicosis[J]. *BMC pulmonary medicine*, 2015, 15(1): 66. DOI: 10.1186 / s12890-015-0061-8.
 - [34] Perotin JM, Jeanfaivre T, Thibout Y, et al. Endoscopic management of idiopathic tracheal stenosis[J]. *Ann Thorac*

- Surg, 2011, 92(1): 297-300. DOI: 10.1016 / j.athoracsur.2011.03.129.
- [35] Debeljak A, Kecelj P. Bronchoscopic removal of foreign bodies in adults: experience with 62 patients from 1974-1998 [J]. *European Respiratory Journal*, 1999, 14(4): 792-795. DOI: 10.1034/j.1399-3003.1999.14d11.x.
 - [36] Tsao TCY, Tsai YH, Lan RS, et al. Treatment for Collapsed Lung in Critically Ill Patients: Selective Intrabronchial Air Insufflation Using the Fiberoptic Bronchoscope[J]. *Chest*, 1990, 97(2): 435-438. DOI:10.1378/chest.97.2.435.
 - [37] Baumgartner F, Sheppard B, de Virgilio C, et al. Tracheal and main bronchial disruptions after blunt chest trauma: presentation and management[J]. *The Annals of thoracic surgery*, 1990, 50(4): 569-574. DOI:10.1016/0003-4975(90)90191-8.
 - [38] Prokakis C, Dougenis D, Koletsis EN, et al. Airway trauma: a review on epidemiology, mechanisms of injury, diagnosis and treatment[J]. *Journal of cardiothoracic surgery*, 2014, 9(1): 117. DOI: 10.1186/1749-8090-9-117.
 - [39] Olson EJ, Utz JP, Prakash UBS. Therapeutic bronchoscopy in broncholithiasis[J]. *American journal of respiratory and critical care medicine*, 1999, 160(3): 766-770. DOI: 10.1164/ajrcm.160.3.9810021.
 - [40] York EL, Lewall DB, Hirji M, et al. Endoscopic diagnosis and treatment of postoperative bronchopleural fistula[J]. *Chest*, 1990, 97(6): 1390-1392. DOI: 10.1378/chest.97.6.1390.
 - [41] Hollaus PH, Lax F, Janakiev D, et al. Endoscopic treatment of postoperative bronchopleural fistula: experience with 45 cases [J]. *The Annals of thoracic surgery*, 1998, 66(3): 923-927. DOI: 10.1016/S0003-4975(98)00589-X.
 - [42] Limper AH, Prakash UB. Tracheobronchial foreign bodies in adults[J]. *Ann Intern Med*, 1990, 112(8): 604-609. DOI: 10.7326/0003-4819-112-8-604.
 - [43] Fang YF, Hsieh MH, Chung FT, et al. Flexible bronchoscopy with multiple modalities for foreign body removal in adults[J]. *PloS one*, 2015, 10(3): e0118993. DOI: 10.1371 / journal.pone.0118993.
 - [44] Lin L, Lv L, Wang Y, et al. The clinical features of foreign body aspiration into the lower airway in geriatric patients[J]. *Clinical interventions in aging*, 2014, 9: 1613. DOI: 10.2147/CIA.S70924.
 - [45] Haponik EF, Shure D. Underutilization of trans-bronchial needle aspiration: experiences of current pulmonary fellows[J]. *Chest*, 1997, 112(1): 251-253. DOI: 10.1378/chest.112.1.251.
 - [46] Sharafkhaneh A, Baaklini W, Gorin AB, et al. Yield of transbronchial needle aspiration in diagnosis of mediastinal lesions[J]. *Chest*, 2003, 124(6): 2131-2135. DOI: 10.1378 / chest.124.6.2131.
 - [47] Prakash UB, Offord KP, Stubbs SE. Bronchoscopy in North America: the ACCP survey[J]. *Chest*, 1991, 100(6): 1668-1675. DOI: 10.1378/chest.100.6.1668.
 - [48] Dweik RA, Mehta AC, Meeker DP, et al. Analysis of the safety of bronchoscopy after recent acute myocardial infarction[J]. *Chest*, 1996, 110(3): 825-828. DOI: 10.1378/chest.110.3.825.
 - [49] Dunagan DP, Burke HL, Aquino SL, et al. Fiberoptic bronchoscopy in coronary care unit patients: indications, safety, and clinical implications[J]. *Chest*, 1998, 114(6): 1660. DOI: 10.1378/chest.114.6.1660.
 - [50] Fleisher LA, Beckman JA, Brown KA, et al. ACC/AHA 2007 guidelines on perioperative cardiovascular evaluation and care for noncardiac surgery[J]. *Circulation*, 2007, 116(2): e418-500. DOI: 10.1016/j.jacc.2007.09.003.
 - [51] Dupree HJ, Lewejohann JC, Gleiss J, et al. Fiberoptic bronchoscopy of intubated patients with life-threatening hemoptysis[J]. *World J Surg*, 2001, 25(1): 104-107. DOI: 10.1007/s002680020366.
 - [52] Khalil A, Soussan M, Mangiapan G, et al. Utility of high-resolution Chest CT scan in the emergency management of haemoptysis in the intensive care unit: severity, localization and aetiology[J]. *Br J Radiol*, 2007, 80(949): 21-25. DOI: 10.1259/bjr/59233312.
 - [53] Weiss SM, Hert RC, Gianola FJ, et al. Complications of fiberoptic bronchoscopy in thrombocytopenic patients[J]. *Chest*, 1993, 104(4): 1025-1028. DOI: 10.1378 / chest.104.4.1025.
 - [54] Diette GB, Wiener CM, White P Jr. The higher risk of bleeding in lung transplant recipients from bronchoscopy is independent of traditional bleeding risks: results of a prospective cohort study[J]. *Chest*, 1999, 115(2): 397-402. DOI:10.1378/chest.115.2.397.
 - [55] Veitch AM, Vanbiervliet G, Gershlick A, et al. Endoscopy in patients on antiplatelet or anticoagulant therapy, including direct oral anticoagulants: British Society of Gastroenterology (BSG) and European Society of Gastrointestinal Endoscopy (ESGE) guidelines[J]. *Endoscopy*, 2016, 65(3): 374-389. DOI: 10.1055/s-0042-102652.
 - [56] Bahhady IJ, Ernst A. Risks of and recommendations for flexible bronchoscopy in pregnancy: a review[J]. *Chest*, 2004, 126(6):1974. DOI:10.1378/chest.126.6.1974.
 - [57] Poi PJ, Chuah SY, Srinivas P, et al. Common fears of patients undergoing bronchoscopy[J]. *Eur Respir J*, 1998, 11(5): 1147-1149. DOI:10.1183/09031936.98.11051147.
 - [58] Lechtzin N, Rubin HR, White P Jr, et al. Patient satisfaction with bronchoscopy[J]. *Am J Respir Crit Care Med*, 2002, 166(10):1326-1331. DOI:10.1164/rccm.200203-2310C.
 - [59] Bernasconi M, Chhaged PN, Muller P, et al. Patients's satisfaction with flexible bronchoscopy in a hospital-based community practice[J]. *Respiration*, 2009, 78(4): 440-445. DOI: 10.1159/000228906.
 - [60] Mitsumune T, Senoh E, Adachi M. Prediction of patient discomfort during fiberoptic bronchoscopy[J]. *Respirology*, 2005, 10(1): 92-96. DOI: 10.1111/j.1440-1843.2005.00642.x.
 - [61] Hadzri H, Azarisman S, Fauzi A, et al. Can a bronchoscopist reliably assess a patient's experience of bronchoscopy? [J]. *RSM Short Rep*, 2010, 1(4): 35. DOI: 10.1258 / shorts.2010.010044.
 - [62] Mitsumune T, Senoh E, Adachi M. Prediction of patient discomfort during fiberoptic bronchoscopy[J]. *Respirology*, 2005, 10(1): 92-96. DOI: 10.1111/j.1440-1843.2005.00642.x.
 - [63] Diette GB, White P Jr, Try P, et al. Quality assessment through patient self-report of symptoms pre-fiberoptic and post-fiberoptic bronchoscopy[J]. *Chest*, 1998, 114: 1446-1453. DOI: 10.1378/chest.114.5.1446.
 - [64] Hadzri H, Azarisman S, Fauzi A, et al. Can a bronchoscopist reliably assess a patients' experience of bronchoscopy? [J]. *JRSM Short Rep*, 2010, 1: 35. DOI: 10.1258 / shorts. 2010. 010044.
 - [65] General Medical Council UK. Consent guidance: patients and doctors making decisions together. 2008. http://www.gmc-uk.org/guidance/ethical_guidance/consent_guidance_index.asp.
 - [66] General Medical Council UK. Consent guidance: patients and doctors making decisions together. 2008. http://www.gmc-uk.org/guidance/ethical_guidance/consent_guidance_index.asp.

- org/guidance/ethical_guidance/consent_guidance_index.asp.
- [67] Meyer KC, Raghu G, Baughman RP, et al. An official American Thoracic Society clinical practice guideline: the clinical utility of bronchoalveolar lavage cellular analysis in interstitial lung disease[J]. *Am J Respir Crit Care Med*, 2012, 185(9): 1004-1014. DOI: 10.1164/rccm.201202-0320st.
 - [68] Rivera MP, Mehta AC. Initial diagnosis of lung cancer[J]. *Chest*, 2007, 132(3 Suppl): 131S-148S. DOI: 10.1378 / chest.07-1357.
 - [69] McLean AN, Semple PA, Franklin DH, et al. The Scottish multi-centre prospective study of bronchoscopy for bronchial carcinoma and suggested audit standards[J]. *Respir Med*, 1998, 92: 1110-1115. DOI: 10.1016/S0954-6111(98)90403-6.
 - [70] Gellert AR, Rudd RM, Sinha G, et al. Fiberoptic bronchoscopy: effect of multiple bronchial biopsies on diagnostic yield in bronchial carcinoma[J]. *Thorax*, 1982, 37: 684-687. DOI: 10.1136/thx.37.9.684.
 - [71] Brady M, Kinn S, Stuart P. Preoperative fasting for adults to prevent perioperative complications[J]. *Cochrane Database Syst Rev*, 2003, 4: CD004423. DOI: 10.1002 / 14651858. CD004423.
 - [72] Molina JA, Lobo CA, Goh HK, et al. Review of studies and guidelines on fasting and procedural sedation at the emergency department[J]. *Int J Evid Based Healthc*, 2010, 8: 75-78. DOI: 10.1111/j.1479-6988.2010.00163.x.
 - [73] British Thoracic Society Bronchoscopy Guidelines Committee. British Thoracic Society guidelines on diagnostic flexible bronchoscopy[J]. *Thorax*, 2001, 56(Suppl 1): i1-21. DOI: 10.1136/thorax.56.suppl_1.i1.
 - [74] Brady M, Kinn S, Stuart P. Preoperative fasting for adults to prevent perioperative complications[J]. *Cochrane Database Syst Rev*, 2003, (4): CD004423. DOI: 10.1002 / 14651858. CD004423.
 - [75] Molina JA, Lobo CA, Goh HK, et al. Review of studies and guidelines on fasting and procedural sedation at the emergency department[J]. *Int J Evid Based Healthc*, 2010, 8: 75-78. DOI: 10.1111/j.1479-6988.2010.00163.x.
 - [76] 中华医学会麻醉学分会. (支)气管镜诊疗镇静/麻醉的专家共识(2014)[J]. 人民卫生出版社, 2014: 613-667.
 - [77] DuRand IA, Blaikley J, Booton R, et al. British Thoracic Society guideline for diagnostic flexible bronchoscopy in adults[J]. *Thorax*, 2013, 68: i1-i144. DOI: 10.1136/thoraxjnl-2013-203629.
 - [78] Malik JA, Gupta D, Agarwal AN, et al. Anticholinergic premedication for flexible bronchoscopy: a randomized, double-blind, placebo-controlled study of atropine and glycopyrrolate[J]. *Chest*, 2009, 136: 347-354. DOI: 10.1378 / chest.08-2924.
 - [79] De Padua AI, de Castro M, Schmidt A, et al. Clonidine as a pre-anesthetic agent for flexible bronchoscopy[J]. *Respir Med*, 2004, 98: 746-751. DOI: 10.1016/j.rmed.2004.01.015.
 - [80] Matot I, Kramer MR. Sedation in outpatient bronchoscopy[J]. *Respir Med*, 2000, 94: 1145-1153. DOI: 10.1053 / rmed. 2000.0926.
 - [81] Williams T, Brooks T, Ward C. The role of atropine premedication in fiberoptic bronchoscopy using intravenous midazolam sedation[J]. *Chest*, 1998, 113: 1394-1398. DOI: 10.1378/chest.113.5.1394.
 - [82] Cowl CT, Prakash UB, Kruger BR. The role of anticholinergics in bronchoscopy. A randomized clinical trial[J]. *Chest*, 2000, 118: 188-192. DOI: 10.1378/chest.118.1.188
 - [83] Kozak EA, Brath LK. Do 'screening' coagulation tests predict bleeding in patients undergoing fiberoptic bronchoscopy with biopsy? [J]. *Chest*, 1994, 106: 703-705. DOI: 10.1378 / chest.106.3.703.
 - [84] Cordasco EM Jr, Mehta AC, Ahmad M. Bronchoscopically induced bleeding. A summary of nine years' Cleveland clinic experience and review of the literature[J]. *Chest*, 1991, 100: 1141-1147. DOI: 10.1378/chest.100.4.1141.
 - [85] Bjortuff O, Brosstad F, Boe J. Bronchoscopy with transbronchial biopsies: measurement of bleeding volume and evaluation of the predictive value of coagulation tests[J]. *Eur Respir J*, 1998, 12: 1025-1027. DOI: 10.1183 / 09031936.98.12051025.
 - [86] Kumar S, Nath A, Singh S, et al. An unusual complication during bronchoscopy: hypotension, global ST segment elevation, and acute severe left ventricular systolic dysfunction [J]. *Respir Care*, 2013, 58(9): e111-e115. DOI: 10.4187 / respcare.02318.
 - [87] Davies L, Mister R, Spence DP, et al. Cardiovascular consequences of fiberoptic bronchoscopy[J]. *Eur Respir J*, 1997, 10 (3): 695-698. DOI: 10.1097/00128594-199707000-00019.
 - [88] Ernst A, Eberhardt R, Wahidi M, et al. Effect of routine clopidogrel use on bleeding complications after transbronchial biopsy in humans[J]. *Chest*, 2006, 129: 734-737. DOI: 10.1378/chest.129.3.734.
 - [89] Veitch AM, Baglin TP, Gershlick SM, et al. Guidelines for the management of anticoagulant and antiplatelet therapy in patients undergoing endoscopic procedures[J]. *Gut*, 2008, 57: 1322-1329. DOI: 10.1136/gut.2007.142497.
 - [90] Gordon HG, Elie AA, Mark C, et al. Managing anticoagulation and antithrombotic in the perioperative period: American College of Chest Physicians (ACCP) guidelines (9th Edition) [J]. *Chest*, 2012, 141(2): 7s-47s. DOI: 10.1378/chest.141283.
 - [91] Peacock AJ, Benson-Mitchell R, Godfrey R. Effect of fiberoptic bronchoscopy on pulmonary function[J]. *Thorax*, 1990, 45: 38-41. DOI: 10.1136/thx.45.1.38.
 - [92] Chechani V. Flexible bronchoscopy in patients with hypercapnia[J]. *J Bronchol*, 2000, 7: 226-232. DOI: 10.1097 / 00128594-200007030-00006.
 - [93] Djukanovic R, Wilson JW, Lai CK, et al. The safety aspects of fiberoptic bronchoscopy, bronchoalveolar lavage, and endobronchial biopsy in asthma[J]. *Am Rev Respir Dis*, 1991, 143(4 Pt 1): 772-777. DOI: 10.1164/ajrccm/143.4_Pt_1.772.
 - [94] Moore WC, Evans MD, Bleeker ER, et al. Safety of investigative bronchoscopy in the Severe Asthma Research Program[J]. *J Allergy Clin Immunol*, 2011, 128: 328-336. DOI: 10.1016/j.jaci.2011.02.042.
 - [95] Humbert M, Robinson DS, Assoufi B, et al. Safety of fiberoptic bronchoscopy in asthmatic and control subjects and effect on asthma control over two weeks[J]. *Thorax*, 1996, 51: 664-669. DOI: 10.1016/S0091-6749(96)80722-6.
 - [96] Elston MJ, Whittaker AJ, Khan LN, et al. Safety of research bronchoscopy, biopsy and bronchoalveolar lavage in asthma [J]. *Eur Respir J*, 2004, 24: 375-377. DOI: 10.1183 / 09031936.04.00063003.
 - [97] Stolz D, Pollak V, Chhajed PN, et al. A randomized, placebo-controlled trial of bronchodilators for bronchoscopy in patients with COPD[J]. *Chest*, 2007, 131: 765-772. DOI: 10.1378/chest.06-2308.
 - [98] Milman N, Faurschou P, Grode G, et al. Pulse oximetry during fiberoptic bronchoscopy in local anaesthesia: frequency of hypoxaemia and effect of oxygen supplementation[J].

- Respiration, 1994, 61: 342-347. DOI: 10.1159/000196366.
- [99] Yildiz P, Ozgul A, Yimaz V. Changes in oxygen saturation in patients undergoing fiberoptic bronchoscopy[J]. Chest, 2002, 121: 1007-1008. DOI: 10.1378/chest.121.3.1007.
- [100] Chechani V. Flexible bronchoscopy in patients with hypercapnia[J]. J Bronchol, 2000, 7: 226-232. DOI: 10.1097/00128594-200007030-00006.
- [101] Maranetra N, Pushpakom R, Bovomkitti S. Oxygen desaturation during fiberoptic bronchoscopy[J]. J Med Assoc Thai, 1990, 73: 258-263.
- [102] Cheng Q, Zhang J, Wang H, et al. Effect of Acute Hypercapnia on Outcomes and Predictive Risk Factors for Complications among Patients Receiving Bronchoscopic Interventions under General Anesthesia[J]. PLoS One, 2015, 10(7): e0130771. DOI: 10.1371/journal.pone.0130771.
- [103] Hirose T, Okuda K, Ishida H, et al. Patient satisfaction with sedation for flexible bronchoscopy[J]. Respirology, 2008, 13: 722-727. DOI: 10.1111/j.1440-1843.2008.01311.x.
- [104] Smyth CM, Stead RJ. Survey of flexible fiberoptic bronchoscopy in the United Kingdom[J]. Eur Respir J, 2002, 19: 458-463. DOI: 10.1183/09031936.02.00103702.
- [105] Maguire GP, Rubinfeld AR, Trembath PW, et al. Patients prefer sedation for fiberoptic bronchoscopy[J]. Respirology, 1998, 3: 81-85. DOI: 10.1111/j.1440-1843.1998.tb00101.x.
- [106] Silvestri GA, Vincent BD, Wahidi MM, et al. A phase 3, randomized, double-blind study to assess the efficacy and safety of fospropofol disodium injection for moderated sedation in patients undergoing flexible bronchoscopy[J]. Chest, 2009, 135: 41-47. DOI: 10.1378/chest.08-0623.
- [107] Hatton MQ, Allen MB, Vathenen AS, et al. Does sedation help in fiberoptic bronchoscopy? [J]. BMJ, 1994, 309: 1206-1207. DOI: 10.1136/bmj.309.6963.1206.
- [108] Gonzalez R, De-La-Rosa-Ramirez I, Maldonado-Hernandez A, et al. Should patients undergoing a bronchoscopy be sedated? [J]. Acta Anaesthesiol Scand, 2003, 47: 411-415. DOI: 10.1034/j.1399-6576.2003.00061.x.
- [109] Putinati S, Ballerin L, Corbetta L, et al. Patient satisfaction with conscious sedation for bronchoscopy[J]. Chest, 1999, 115: 1437-1440. DOI: 10.1378/chest.115.5.1437.
- [110] Hirose T, Okuda K, Ishida H, et al. Patient satisfaction with sedation for flexible bronchoscopy[J]. Respirology, 2008, 13: 722-727. DOI: 10.1111/j.1440-1843.2008.01311.x.
- [111] Reducing risk of overdose with midazolam injection in adults. NPSA / 2008 / RRR011. 2008. <http://www.nrls.npsa.uk/resources/?entryid45=59896> (accessed Feb 2019).
- [112] Chernik DA, Gillings D, Laine H, et al. Validity and reliability of the Observer's Assessment of Alertness/Sedation Scale; study with intravenous midazolam[J]. J Clin Psychopharmacol, 1990, 10: 244-251. DOI: 10.1097/00004714-199008000-00003.
- [113] Clarkson K, Power CK, O'Connell F, et al. A comparative evaluation of Propofol and midazolam as sedative agents in fiberoptic bronchoscopy[J]. Chest, 1993, 104: 1029-1031. DOI: 10.1378/chest.104.4.1029.
- [114] Clark G, Licker M, Younossian AB, et al. Titrated sedation with Propofol or midazolam for flexible bronchoscopy: a randomized trial[J]. Eur Respir J, 2009, 34: 1277-1283. DOI: 10.1183/09031936.00142108.
- [115] Crawford M, Pollock J, Anderson K, et al. Comparison of midazolam with Propofol for sedation in outpatient bronchoscopy[J]. Br J Anaesth, 1993, 70: 419-422. DOI: 10.1093/bja/70.4.419.
- [116] Houghton CM, Raghuram A, Sullivan PJ, et al. Pre-medication for bronchoscopy: randomized double blind trial comparing alfentanil with midazolam[J]. Respir Med, 2004, 98: 1102-1107. DOI: 10.1016/j.rmed.2004.03.023.
- [117] Lo YL, Lin TY, Fang YF, et al. Feasibility of bispectral index-guided Propofol infusion for flexible bronchoscopy sedation: a randomized controlled trial[J]. PLoS ONE, 2011, 6: e27769. DOI: 10.1371/journal.pone.0027769.
- [118] Stolz D, Chhajed PN, Leuppi JD, et al. Cough suppression during flexible bronchoscopy using combined sedation with midazolam and hydrocodone: a randomized, double blind, placebo controlled trial[J]. Thorax, 2004, 59: 773-776. DOI: 10.1136/thx.2003.019836.
- [119] Greig JH, Cooper SM, Kasimbazi HJ, et al. Sedation for fiberoptic bronchoscopy[J]. Respir Med, 1995, 89: 53-56. DOI: 10.1016/0954-6111(95)90071-3.
- [120] Gray AJG, Bell GD. Elderly patients vulnerable because of excessive doses of sedatives. National Confidential Enquiry into Patient Outcome and Death(NCEPOD). <http://www.ncepod.org.uk/pdf/current/NPSA%20sedation%20article.pdf> (accessed Feb 2019).
- [121] 邓丽云, 赵真英, 郭晋铎, 等. 国产咪唑啉仑用于静脉诱导的药代动力学[J]. 中国药物与临床, 2008, 8(9): 741-742. DOI: 10.3969/j.issn.1671-2560.2008.09.029.
- [122] 李爱君, 滕海风, 张文强, 等. 咪达唑仑和安定在气管镜检查中应用的比较[J]. 中华麻醉学杂志, 2004, 24(9): 708-709. DOI: 10.3760/j.issn:0254-1416.2004.09.022.
- [123] 罗朝志, 岳云, 黄雄庆, 等. 局麻患者不同剂量咪达唑仑复合芬太尼镇静效果的比较——随机、双盲、多中心研究[J]. 中华麻醉学杂志, 2008, 28(3): 284-285. DOI: 10.3321/j.issn:0254-1416.2008.03.020.
- [124] 宋瑾瑾, 俞万钧. 咪达唑仑清醒镇静在气管镜检查中的应用[J]. 临床肺科杂志, 2008, 13(11): 1511. DOI: 10.3969/j.issn.1009-6663.2008.11.073.
- [125] Schlatter L, Pflimlin E, Fehrke B, et al. Propofol versus Propofol plus hydrocodone for flexible bronchoscopy: a randomized study[J]. Eur Respir J, 2011, 38: 529-537. DOI: 10.1183/09031936.00121610.
- [126] Stolz D, Kurer G, Meyer A, et al. Propofol versus combined sedation in flexible bronchoscopy: a randomized non-inferiority trial[J]. Eur Respir J, 2009, 34: 1024-1030. DOI: 10.1183/09031936.00180808.
- [127] Dreher M, Ekkernkamp E, Storre JH, et al. Sedation during flexible bronchoscopy in patients with pre-existing respiratory failure: midazolam versus midazolam plus alfentanil[J]. Respiration, 2010, 79: 307-314. DOI: 10.1159/000267227.
- [128] Yuan F, Fu HG, Yang PJ, et al. Dexmedetomidine-fentanyl versus Propofol-fentanyl in flexible bronchoscopy: A randomized study[J]. Exp Ther Med, 2016, 12(1): 506-512. DOI: 10.3892/etm.2016.3274.
- [129] Goneppanavar U, Magazine R, Periyadka B, et al. Intravenous dexmedetomidine provides superior patient comfort and tolerance compared to intravenous midazolam in patients undergoing flexible bronchoscopy[J]. Pulm Med, 2015, 2015: 727530. DOI: 10.1155/2015/727530.
- [130] Mondal S, Ghosh S, Bhattacharya S, et al. Comparison between dexmedetomidine and fentanyl on intubation conditions during awake fiberoptic bronchoscopy: A randomized double-blind prospective study[J]. J Anaesthesiol Clin Pharmacol, 2015, 31(2): 212-216. DOI: 10.4103 /

- 0970-9185.155151.
- [131] Gao Y, Kang K, Liu H, et al. Effect of dexmedetomidine and midazolam for flexible fiberoptic bronchoscopy in intensive care unit patients: A retrospective study[J]. *Kaijiang Yu Medicine*, 2017, 96(25): e7090. DOI: 10.1097 / MD.0000000000007090.
 - [132] 中华医学会麻醉学分会. 右美托咪定临床应用指导意见 (2013)[J]. *中华麻醉学杂志*, 2013, 33(10): 1165-1167. DOI: 10.3760/cma.j.issn.0254-1416.2013.10.001.
 - [133] Antoniadis N, Worsnop C. Topical lidocaine through the bronchoscope reduces cough rate during bronchoscopy[J]. *Respirology*, 2009, 14: 873-876. DOI: 10.1111 / j. 1440-1843. 2009.01587.x.
 - [134] Guay J. Methemoglobinemia related to local anesthetics: a summary of 242 episodes[J]. *Anesth Analg*, 2009, 108: 837-845. DOI: 10.1213/ane.0b013e318187c4b1.
 - [135] Randell T, Yli-Hankala A, Valli H, et al. Topical anaesthesia of the nasal mucosa for fiberoptic airway endoscopy[J]. *Br J Anaesth*, 1992, 68: 164-167. DOI: 10.1093/bja/68.2.164.
 - [136] Webb AR, Woodhead MA, Dalton HR, et al. Topical nasal anaesthesia for fiberoptic bronchoscopy: patients' preference for lignocaine gel[J]. *Thorax*, 1989, 44: 674-675. DOI: 10.1136/thx.44.8.674.
 - [137] Zainudin BM, Rafia MH, Sufarlan AW. Topical nasal anaesthesia for fiberoptic bronchoscopy: lignocaine spray or gel? [J]. *Singapore Med J*, 1993, 34: 148-149. DOI: 10.1136/ thx.45.1.79.
 - [138] Keane D, McNicholas WT. Comparison of nebulized and sprayed topical anaesthesia for fiberoptic bronchoscopy[J]. *Eur Respir J*, 1992, 5: 1123-1125.
 - [139] Webb J. Local anaesthesia for fiberoptic bronchoscopy—where are we now? [J]. *Respir Med*, 1990, 84: 349-350. DOI: 10.1016/S0954-6111(08)80067-4.
 - [140] Isaac PA, Barry JE, Vaughan RS, et al. A jet nebulizer for delivery of topical anesthesia to the respiratory tract. A comparison with cricothyroid puncture and direct spraying for fiberoptic bronchoscopy[J]. *Anaesthesia*, 1990, 45: 46-48.
 - [141] Stolz D, Chhajed PN, Leuppi J, et al. Nebulized lidocaine for flexible bronchoscopy: a randomized, double-blind, placebo-controlled trial[J]. *Chest*, 2005, 128: 1756-1760. DOI: 10.1378/chest.128.3.1756.
 - [142] Hasmoni MH, Rani MFA, Harun R, et al. randomized-controlled trial to study the equivalence of 1% versus 2% lignocaine in cough suppression and satisfaction during bronchoscopy[J]. *J Bronchol Interv Pulmonol*, 2008, 15: 78-82. DOI: 10.1097/lbr.0b013e31816b653c.
 - [143] Mainland PA, Kong AS, Chung DC, et al. Absorption of lidocaine during aspiration anesthesia of the airway[J]. *J Clin Anesth*, 2001, 13: 440-446. DOI: 10.1016 / S0952-8180(01) 00298-7.
 - [144] Xue FS, Liu HP, He N, et al. Spray-as-you-go airway topical anesthesia in patients with a difficult airway: a randomized, double-blind comparison of 2% and 4% lidocaine[J]. *Anesth Analg*, 2009, 108: 536-543. DOI: 10.1213/ane.0b013e31818f1665.
 - [145] 程玉生, 陆志伟. 利多卡因对支气管镜检查的麻醉效果评价[J]. *中华肺部疾病杂志(电子版)*, 2013, 6(1): 54-56. DOI: 10.3877/cma.j.issn.1674-6902.2013.01.011.
 - [146] Efthimiou J, Higenbottam T, Holt D, et al. Plasma concentrations of lignocaine during fiberoptic bronchoscopy [J]. *Thorax*, 1982, 37: 68-71. DOI: 10.1136/thx.37.1.68.
 - [147] Milman N, Laub M, Munch EP, et al. Serum concentrations of lignocaine and its metabolite monoethylglycinexylidide during fibre-optic bronchoscopy in local anaesthesia[J]. *Respir Med*, 1998, 92: 40-43. DOI: 10.1016/S0954-6111(98)90030-0.
 - [148] Frey WC, Emmons EE, Morris MJ. Safety of high dose lidocaine in flexible bronchoscopy[J]. *J Bronchol*, 2008, 15: 33-37. DOI: 10.1097/lbr.0b013e3181641b8e.
 - [149] Langmack EL, Martin FJ, Pak J, et al. Serum lidocaine concentrations in asthmatics undergoing research bronchoscopy[J]. *Chest*, 2000, 117: 1055-1060. DOI: 10.1378/ chest.117.4.1055.
 - [150] Ameer B, Brulingame MB, Harman EM. Systemic absorption of topical lidocaine in elderly and young adults undergoing bronchoscopy[J]. *Pharmacotherapy*, 1989, 9: 74-81. DOI: 10.1002/j.1875-9114.1989.tb04106.x.
 - [151] Markou Nk, Kanakaki MC, Boutzouka E, et al. Fluctuations in gas exchange and cardiovascular parameters during flexible bronchoscopy[J]. *J Bronchol*, 1999, 6: 241-246. DOI: 10.1097/ 00128594-199910000-00003.
 - [152] Davies L, Mister R, Spence DP, et al. Cardiovascular consequences of fiberoptic bronchoscopy[J]. *Eur Respir J*, 1997, 10: 695-698. DOI: 10.1097/00128594-199707000-00019.
 - [153] Ouellette DR, Diaz J. Elevation of the double product during flexible bronchoscopy: effects of uncontrolled hypertension and the use of beta-blockade[J]. *J Bronchol*, 2008, 15: 73-77. DOI: 10.1097/lbr.0b013e318169e2e8.
 - [154] Sharma GD, Bansal SK, Kashyap S, et al. Effect of fiberoptic bronchoscopy on arterial blood gases and cardiac rhythm at a moderate altitude of 2250 meters[J]. *J Assoc Physicians India*, 1999, 47(11): 1056-1059.
 - [155] Van Zwam JP, Kapteijns EFG, Lhey S, et al. Flexible bronchoscopy in supine or sitting position: a randomized prospective analysis of safety and patient comfort[J]. *J Bronchol*, 2010, 17: 29-32. DOI: 10.1097 / LBR. 0b013e3181cc3a75.
 - [156] Fang WF, Chen YC, Chung YH, et al. Predictors of oxygen desaturation in patients undergoing diagnostic bronchoscopy [J]. *Chang Gung Med J*, 2006, 29: 306-312.
 - [157] Jones AM, O'Driscoll R. Do all patients require supplemental oxygen during flexible bronchoscopy? [J]. *Chest*, 2001, 119: 1906-1909. DOI: 10.1378/chest.119.6.1906.
 - [158] Schiffman PL, Westlake RE, Fourie JA, et al. Arterial oxygen saturation and cardiac rhythm during transoral fiberoptic bronchoscopy[J]. *J Med Soc N J*, 1982, 79: 723-726.
 - [159] Milman N, Faurschou P, Munch EP, et al. Transbronchial lung biopsy through the fiberoptic bronchoscope. Result and complications in 452 examinations[J]. *Respir Med*, 1994, 88: 749-753. DOI: 10.1016/S0954-6111(05)80197-0.
 - [160] Izbicik G, Shitrit D, Yarmolovsky A, et al. Indications, diagnostic yields and complications of transbronchial biopsy over 5 years in the State of Qatar[J]. *Saudi Med J*, 2005, 26: 641-645. DOI: 10.1016/j.revmed.2004.11.008.
 - [161] Ahmad M, Livingston DR, Golish JA, et al. The safety of outpatient transbronchial biopsy[J]. *Chest*, 1986, 90: 403-405. DOI: 10.1378/chest.90.3.403.
 - [162] Milam MC, Evins AE, Sahn SA. Immediate chest roentgenography following fiberoptic bronchoscopy[J]. *Chest*, 1989, 96: 477-479. DOI: 10.1378/chest.96.3.477.
 - [163] Colt HG, Matsuo T. Hospital charges attributable to bronchoscopy-related complications in outpatient[J]. *Respiration*, 2001, 68: 67-72. DOI: 10.1159/000050465.
 - [164] Sharif-Kashani B, Shahabi P, Behzadnia N, et al. Incidence of

- fever and bacteremia following flexible fiberoptic bronchoscopy: a prospective study[J]. *Acta Med Iran*, 2010, 48(6):385-388.
- [165] Centre for Clinical Practice at NICE. Prophylaxis against infective endocarditis(CG64). London: National Institute for Health and Clinical Excellence(NICE), 2008.
- [166] Yigla M, Oren I, Bentur L, et al. Incidence of bacteraemia following fiberoptic bronchoscopy[J]. *Eur Respir J*, 1999, 14: 789-791. DOI: 10.1034/j.1399-3003.1999.14d10.x.
- [167] Haynes J, Greenston MA. Fiberoptic bronchoscopy and the use of antibiotic prophylaxis[J]. *Br Med J Clin Res ed*, 1987, 294:1199. DOI: 10.1136/bmj.294.6581.1199.
- [168] Park JS, Lee CH, Yim JJ, et al. Impact of antibiotic prophylaxis on postbronchoscopy fever: a randomized controlled study[J]. *Int J Tuberc Lung Dis*, 2011, 15: 528-535. DOI:10.5588/ijtld.10.0386.
- [169] Huang JCT, Bassett MA, Levin D, et al. Acute phase reaction in healthy volunteers after bronchoscopy with lavage[J]. *Chest*, 2006, 129: 1565-1569. DOI: 10.1378/chest.129.6.1565.
- [170] Um SW, Choi CM, Lee CT, et al. Prospective analysis of clinical characteristics and risk factors of postbronchoscopy fever[J]. *Chest*, 2004, 125: 945-952. DOI: 10.1378 / chest.125.3.945.
- [171] 中华医学会呼吸病学分会,中国肺癌防治联盟. 肺癌小样本取材相关问题的中国专家共识[J]. *中华内科杂志*, 2016, 55(5):406-413. DOI: 10.3760/cma.j.issn.0578-1426.2016.05.017.
- [172] Mazzone P, Jain P, Arroliga AC, et al. Bronchoscopy and needle biopsy techniques for diagnosis and staging of lung cancer[J]. *Clin Chest Med*, 2002, 23(1): 137-158. DOI: 10.1016/S0272-5231(03)00065-0.
- [173] Gellert AR, Rudd RM, Sinha G, et al. Fiberoptic bronchoscopy: effect of multiple bronchial biopsies on diagnostic yield in bronchial carcinoma[J]. *Thorax*, 1982, 37(9): 684-687. DOI: 10.1136/thx.37.9.684.
- [174] Rivera MP, Mehta AC, Momen M. Establishing the diagnosis of lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines[J]. *Chest*, 2013, 143(5Suppl): e142S-65S. DOI: 10.1378/chest.12-2353.
- [175] Rivera MP, Mehta AC. Initial diagnosis of lung cancer[J]. *Chest*, 2007, 132(3suppl): 131S-48S. DOI: 10.1378 / chest.07-1357.
- [176] McLean AN, Semple PA, Franklin DH, et al. The Scottish multi-centre prospective study of bronchoscopy for bronchial carcinoma and suggested audit standards[J]. *Respir Med*, 1998, 92: 1110-1115. DOI: 10.1016/S0954-6111(98)90403-6.
- [177] Ernst A, Herth FJF. Principles and practice of interventional pulmonology[J]. New York: Springer, 2013: 165-176. DOI: 10.1007/978-1-4614-4292-9.
- [178] Bodh A, Kaushal V, Kashyap S, et al. Cytohistological correlation in diagnosis of lung tumors by using fiberoptic bronchoscopy: study of 200 cases[J]. *Indian J Pathol Microbiol*, 2013, 56(2): 84-88. DOI: 10.4103 / 0377-4929.118661.
- [179] Popovich J Jr, Kvale PA, Eichenhorn MS, et al. Diagnostic accuracy of multiple biopsies from flexible fiberoptic bronchoscopy. A comparison of central versus peripheral carcinoma[J]. *Am Rev Respir Dis*, 1982, 125(5):521-523. DOI: 10.1164/arrd.1982.125.5.521.
- [180] Slade MG, Rahman NM, Stanton AE, et al. Improving standards in flexible bronchoscopy for lung cancer[J]. *Eur Respir J*, 2011, 37: 895-901. DOI: 10.1183/09031936.00097110.
- [181] Hetzel J, Eberhardt R, Herth FJF, et al. Cryobiopsy increases the diagnostic yield of endobronchial biopsy: a multicenter trial[J]. *Eur Respir J*, 2012, 39: 685-690. DOI: 10.1183 / 09031936.00033011.
- [182] Van der Drift MA, van der Wilt GJ, Thunnissen FBJM, et al. A prospective study of the timing and cost-effectiveness of bronchial washing during bronchoscopy for pulmonary malignant tumors[J]. *Chest*, 2005, 128:394-400. DOI: 10.1378/chest.128.1.394.
- [183] Chen W, Gao X, Tian Q, et al. A comparison of autofluorescence bronchoscopy and white light bronchoscopy in detection of lung cancer and preneoplastic lesions: a meta-analysis[J]. *Lung Cancer*, 2011, 73(2): 183-188. DOI: 10.1016/j.lungcan.2010.12.002.
- [184] Herth FJ, Eberhardt R, Anantham D, et al. Narrow-band imaging bronchoscopy increases the specificity of bronchoscopic early lung cancer detection[J]. *J Thorac Oncol*, 2009, 4(9): 1060-1065. DOI: 10.1097 / JTO.0b013e3181b24100.
- [185] Harrow EM, Abi-Saleh W, Blum J, et al. The utility of transbronchial needle aspiration in the staging of bronchogenic carcinoma[J]. *Am J Respir Crit Care Med*, 2000, 161(2 Pt 1):601-607. DOI: 10.1164/ajrcm.161.2.9902040.
- [186] Salathe M, Soler M, Bolliger CT, et al. Transbronchial needle aspiration in routine fiberoptic bronchoscopy[J]. *Respiration*, 1992, 59:5-8.
- [187] Trisolini R, Cancellieri A, Tinelli C, et al. Rapid on-site evaluation of transbronchial aspirates in the diagnosis of hilar and mediastinal adenopathy: a randomized trial[J]. *Chest*, 2011, 139(2): 395-401. DOI: 10.1378/chest.10-1521.
- [188] Oki M, Saka H, Kitagawa C, et al. Rapid on-site cytologic evaluation during endobronchial ultrasound-guided transbronchial needle aspiration for diagnosing lung cancer: a randomized study[J]. *Respiration*, 2013, 85(6): 486-492. DOI: 10.1159/000346987.
- [189] Griffin AC, Schwartz LE, Baloch ZW. Utility of on-site evaluation of endobronchial ultrasound-guided transbronchial needle aspiration specimens[J]. *Cytojournal*, 2011, 8: 20. DOI: 10.4103/1742-6413.90081.
- [190] Murakami Y, Oki M, Saka H, et al. Endobronchial ultrasound-guided transbronchial needle aspiration in the diagnosis of small cell lung cancer[J]. *Respir Investig*, 2014, 52(3): 173-178. DOI: 10.1016/j.resinv.2013.11.004.
- [191] Eapen GA, Shah AM, Lei X, et al. Complications, consequences, and practice patterns of endobronchial ultrasound-guided transbronchial needle aspiration: Results of the AQuIRE registry[J]. *Chest*, 2013, 143(4): 1044-1053. DOI: 10.1378/chest.12-0350.
- [192] Nakajima T, Yasufuku K, Saegusa F, et al. Rapid on-site cytologic evaluation during endobronchial ultrasound-guided transbronchial needle aspiration for nodal staging in patients with lung cancer[J]. *Am Thorac Surg*, 2013, 95(5): 1695-1699. DOI: 10.1016/j.athoracsur.2012.09.074.
- [193] Trisolini R, Cancellieri A, Tinelli C, et al. Randomized trial of endobronchial ultrasound-guided transbronchial needle aspiration with and without rapid on-site evaluation for lung cancer genotyping[J]. *Chest*, 2015, 148(6): 1430-1437. DOI: 10.1378/chest.15-0583.
- [194] Wang Memoli JS, Nietert PJ, Silvestri GA. Meta-analysis of guided bronchoscopy for the evaluation of the pulmonary

- nodule[J]. *Chest*, 2012, 142(2): 385-393. DOI: 10.1378 / chest.11-1764.
- [195] Boonsamrungsuk V, Raweelert P, Juthakarn S. Endobronchial ultrasound plus fluoroscopy versus fluoroscopy-guided bronchoscopy: a comparison of diagnostic yields in peripheral pulmonary lesions[J]. *Lung*, 2012, 190(2): 233-237. DOI: 10.1007/s00408-011-9359-3.
- [196] Ishida T, Asano F, Yamazaki K, et al. Virtual bronchoscopic navigation combined with endobronchial ultrasound to diagnose small peripheral pulmonary lesions: a randomized trial[J]. *Thorax*, 2011, 66(12): 1072-1077. DOI: 10.1136 / thx.2010.145490.
- [197] 唐纯丽, 罗为展, 钟长镐, 等. 径向超声联合虚拟导航引导肺活检对肺外周结节的诊断价值[J]. *中华结核和呼吸杂志*, 2016, (1): 38-40. DOI: 10.3760/cma.j.issn. 1001-0939. 2016. 01. 011.
- [198] Oki M, Saka H, Kitagawa C, et al. Endobronchial ultrasound-guided transbronchial biopsy using novel thin bronchoscope for diagnosis of peripheral pulmonary lesions[J]. *J Thorac Oncol*, 2009, 4(10): 1274-1277. DOI: 10.1097 / JTO.0b013e3181b623e1.
- [199] 张素娟, 周军, 张秋娣, 等. 细支气管镜下超声引导联合测量技术诊断肺周围性病变[J]. *中华结核和呼吸杂志*, 2015, 38 (8): 566-569. DOI: 10.3760/cma.j.issn. 1001-0939. 2015.08.003.
- [200] Zaric B, Perin B, Stojic V, et al. Detection of premalignant bronchial lesions can be significantly improved by combination of advanced bronchoscopic imaging techniques [J]. *Ann Thorac Med*, 2013, 8(2): 93-98. DOI: 10.4103 / 1817-1737.109820.
- [201] Bradley B, Branley HM, Egan JJ, et al. Interstitial lung disease guideline: the British Thoracic Society in collaboration with the Thoracic Society of Australia and New Zealand and the Irish Thoracic Society[J]. *Thorax*, 2008, 63(Suppl 5): v1-58. DOI: 10.1136/thx.2008.101691.
- [202] Milman N, Graudal N, Jacobsen GK. Bronchoalveolar lavage in radiologically detected diffuse lung disease. Diagnostic value of total and differential cell count in a series of 130 patients[J]. *APMIS*, 1995, 103: 764-768.
- [203] Mitchell DM, Emerson CJ, Collins JV, et al. Transbronchial lung biopsy with the fiberoptic bronchoscope: analysis of results in 433 patients[J]. *Br J Dis Chest*, 1981, 75: 258-262.
- [204] Anders GT, Johnson JE, Bush BA, et al. Transbronchial biopsy without fluoroscopy. A seven-year perspective[J]. *Chest*, 1988, 94: 557-560.
- [205] Rhee CK, Kang HH, Kang JY, et al. Diagnostic yield of flexible bronchoscopy without fluoroscopic guidance in evaluating peripheral lung lesions[J]. *J Bronchol*, 2010, 17: 317-322. DOI: 10.1097/LBR.0b013e3181f552a5.
- [206] De Fenoyl O, Capron F, Lebeau B, et al. Transbronchial biopsy without fluoroscopy: a five year experience in outpatients[J]. *Thorax*, 1989, 44: 956-959.
- [207] Descombes E, Gardiol D, Leuenberger P. Transbronchial lung biopsy: an analysis of 530 cases with reference to the number of samples[J]. *Monaldi Arch Chest Dis*, 1997, 52: 324-329.
- [208] Mikolasch TA, Porter JC. Transbronchial lung cryobiopsy in the diagnosis of fibrotic interstitial lung diseases[J]. *PLoS One*. 2014, 28,9(2): e86716. DOI: 10.1371/journal.pone.0086716.
- [209] Dhooria S, Mehta R, Srinivasan A, et al. A multicenter study on the safety and efficacy of different methods for obtaining transbronchial lung cryobiopsy in diffuse lung diseases[J]. *Clin Respir J*, 2018, 12(4): 1711-1720. DOI: 10.1111 / crj.12734. DOI: 10.1111/crj.12734.
- [210] Johansson KA, Marcoux VA, Ronksley PE, et al. Diagnostic Yield and Complications of Transbronchial Lung Cryobiopsy for Interstitial Lung Disease[J]. A Systematic Review and Meta analysis. *Ann Am Thorac Soc*, 2016, 13(10): 1828-1838. DOI: 10.1513/AnnalsATS.201606-461SR.
- [211] 江瑾玥, 郭述良, 李一诗. 经支气管冷冻肺活检技术进展 [J]. *中华结核和呼吸杂志*, 2017, 40(8): 619-622. DOI: 10.3760 / cma.j.issn.1001-0939.2017.08.015.
- [212] Pajares V, Puzo C, Castillo D, et al. Diagnostic yield of transbronchial cryobiopsy in interstitial lung disease: a randomized trial[J]. *Respirology*, 2014, 19 (6): 900-906. DOI: 10.1111/resp.12322.
- [213] Hetzel J, Maldonado F, Ravaglia C, et al. Transbronchial cryopsies for the diagnosis of diffuse parenchymal lung diseases: expert statement from the cryobiopsy working group on safety and utility and a call for standardization of the procedure[J]. *Respiration*, 2018, 95(3): 188-200. DOI: 10.1159 / 000484055.
- [214] 王广发, 陈茂森, 李桂莲, 等. 纤维支气管镜在结节病的所见及其诊断价值[J]. *中国内镜杂志*, 1999, (4): 16-17.
- [215] 张红, 王广发, 章巍, 等. 超声引导下经支气管镜针吸活检对结节病的诊断价值[J]. *中华结核和呼吸杂志*, 2014, 37(10): 774-777. DOI: 10.3760/cma.j.issn.1001-0939.2014.10.018.
- [216] Reynolds HY. Present status of bronchoalveolar lavage in interstitial lung disease[J]. *Curr Opin Pulm Med*, 2009, 15: 479-485. DOI: 10.1097/MCP.0b013e32832d03ef.
- [217] Leonard C, Tormey VJ, O'Keane C, et al. Bronchoscopic diagnosis of sarcoidosis[J]. *Eur Respir J*, 1997, 10(12): 2722-2724.
- [218] Puar HS, Young RC Jr, Armstrong EM. Bronchial and transbronchial lung biopsy without fluoroscopy in sarcoidosis [J]. *Chest*, 1985, 87(3): 303-306.
- [219] Shorr AF, Torrington KG, Hnatiuk OW. Endobronchial biopsy for sarcoidosis: a prospective study[J]. *Chest*, 2001, 120(1): 109-114.
- [220] Rohatgi PK, Kuzmowych TV, Delaney MD. Indications for transbronchial lung biopsy in the diagnosis of intrathoracic sarcoidosis[J]. *Respiration*, 1981, 42: 155-160. DOI: 10.1159 / 000194422.
- [221] Navani N, Booth HL, Kocjan G, et al. Combination of endobronchial ultrasound-guided transbronchial needle aspiration with standard bronchoscopic techniques for the diagnosis of stage I and stage II pulmonary sarcoidosis[J]. *Respirology*, 2011, 16(3): 467-472. DOI: 10.1111 / j.1440-1843.2011.01933.x.
- [222] Abramson MJ, Stone CA, Holmes PW, et al. The role of bronchoalveolar lavage in the diagnosis of suspected opportunistic pneumonia[J]. *Aust M Z J med*, 1987, 17(4): 407-412.
- [223] Rano A, Agusti C, Jimenez P, et al. Pulmonary infiltrates in non-HIV immunocompromised patients: a diagnostic approach using non-invasive and bronchoscopic procedures[J]. *Thorax*, 2001, 56(12): 379-387.
- [224] Orenstein M, Webber CA, Cash M, et al. Value of bronchoalveolar lavage in the diagnosis of pulmonary infection in acquired immune deficiency syndrome[J]. *Thorax*, 1986, 41 (5): 345-349.
- [225] Gracia JD, Miravittles M, Mayordomo C, et al. Empiric treatments impair the diagnostic yield of BAL in HIV-positive patients[J]. *Chest*, 1997, 111(5): 1180-1186.

- [226] Kibiki GS, Mulder B, van der Ven AJ, et al. Laboratory diagnosis of pulmonary tuberculosis in TB and HIV endemic settings and the contribution of real time PCR for M. Tuberculosis in bronchoalveolar lavage fluid[J]. Trop Med Int Health, 2007, 12: 1210-1217. DOI: 10.1111 / j.1365-3156.2007.01907.x.
- [227] Baughman RP, Dohn MN, Loudon RG, et al. Bronchoscopy with bronchoalveolar lavage in tuberculosis and fungal infections[J]. Chest, 1991,99(1): 92-97.
- [228] Salzman SH, Schindel ML, Aranda CP, et al. The role of bronchoscopy in the diagnosis of pulmonary tuberculosis in patients at risk for HIV infection[J]. Chest, 1992, 102(1): 143-146.
- [229] Kennedy DJ, Lewis WP, Barnes PF. Yield of bronchoscopy for the diagnosis of tuberculosis in patients with human immunodeficiency virus infection[J]. Chest, 1992, 102(4): 1040-1044.
- [230] Miro AM, Gibilara E, Powell S, et al. The role of fiberoptic bronchoscopy for diagnosis of pulmonary tuberculosis in patients at risk for AIDS[J]. Chest, 1992,101(5): 1211-1214.
- [231] 王凌航, 毛羽, 赵红心, 等. 支气管镜检查对艾滋病患者肺部感染病原学的诊断价值[J]. 中华结核和呼吸杂志, 2010,33(10):727-729. DOI: 10.3760/cma.j.issn. 1001-0939. 2010.10.004.
- [232] 牟向东, 高莉, 王仁贵, 等. 非艾滋病免疫抑制宿主肺孢子菌肺炎影像学特点及其与预后的关系[J]. 中华医学杂志, 2012, 92(38): 2703-2706. DOI: 10.3760 / cma. j. issn. 0376-2491.2012.38.011.
- [233] Yoshimoto E, Konishi M, Takahashi K, et al. BALF findings in HIV-infected patients with Pneumocystis Carinii Pneumonia [J]. Journal of the Japan Society for Bronchology, 2001, 23: 527-531.
- [234] Yu Q, Jia P, Su L, et al. Outcomes and prognostic factors of non-HIV patients with pneumocystis jirovecii pneumonia and pulmonary CMV co-infection: A Retrospective Cohort Study [J]. BMC Infect Dis, 2017, 17(1): 392. DOI: 10.1186 / s12879-017-2492-8.
- [235] Beam E, Germer JJ, Lahr B, et al. Cytomegalovirus (CMV) DNA quantification in bronchoalveolar lavage fluid of immunocompromised patient with CMV pneumonia[J]. Clinical Transplantation, 2018, 32(1): e13149. DOI: 10.1111/ctr.13149.
- [236] Henke-Gendo C, Ganzenmueller T, Kluba J, et al. Improved quantitative PCR protocols for adenovirus and CMV with an internal inhibition control system and automated nucleic acid isolation[J]. J Med Virol, 2012,84(6):890-896. DOI: 10.1002/jmv.23285.
- [237] Jouneau S, Poineuf JS, Minjolle S, et al. Which patients should be tested for viruses on bronchoalveolar lavage fluid? [J]. European Journal of Clinical Microbiology & Infectious Diseases, 2013,32(5): 671-677. DOI:10.1007/s10096-01201791-7.
- [238] Dasgupta KS, Mundada PS, Soni N. Diagnostic role of fiberoptic bronchoscopy in pulmonary tuberculosis[J]. Indian J Otolaryngol Head Neck Surg, 2000, 52: 347-349.
- [239] de Gracia J, Curull V, Vidal R, et al. Diagnostic value of bronchoalveolar lavage in suspected pulmonary tuberculosis [J]. Chest, 1988,93: 329-332.
- [240] Altaf Bachh A, Gupta R, Haq I, et al. Diagnosing sputum / smear-negative pulmonary tuberculosis: does fibre-optic bronchoscopy play a significant role? [J]Lung India, 2010, 27: 58-62. DOI: 10.4103/0970-2113.63607.
- [241] Liam Ck, Chen YC, Yap SF, et al. Detection of Mycobacterium tuberculosis in bronchoalveolar lavage from patients with sputum smear-negative pulmonary tuberculosis using a polymerase chain reaction assay[J]. Respirology, 1998, 3: 125-129.
- [242] Tueller C, Chhajed PN, Buitrago-Tellez C, et al. Value of smear and PCR in bronchoalveolar lavage fluid in culture positive pulmonary tuberculosis[J]. Eur Respir J, 2005, 26: 767-772. DOI: 10.1183/09031936.05.00046105.
- [243] Jafari C, Thijsen S, Sotgiu G, et al. Bronchoalveolar lavage enzyme-linked immunospot for a rapid diagnosis of tuberculosis: a Tuberculosis Network European Trials group study[J]. Am J Respir Crit Care Med, 2009, 180: 666-673. DOI: 10.1164/rccm.200904-0557OC.
- [244] Heurlin N, Elvin K, Lidman C, et al. Fiberoptic bronchoscopy and sputum examination for diagnosis of pulmonary disease in AIDS patients in Stockholm[J]. Scand J Infect Dis, 1990, 22: 659-664.
- [245] Charoenratanakul S, Dejsomritrutai W, Chaiprasert A. Diagnostic role of fiberoptic bronchoscopy in suspected smear negative pulmonary tuberculosis[J]. Respir Med, 1995, 89: 621-623.
- [246] Jaiswal AK, Kulpati DD, Jain NK, et al. Role of bronchoacopy in the early diagnosis of suspected smear negative cases of pulmonary tuberculosis[J]. Ind J Tub, 1989,36: 233.
- [247] De Gracia J, Curull V, Vidal R, et al. Diagnostic value of bronchoalveolar lavage in suspected pulmonary tuberculosis [J]. Chest, 1988,93(2): 329-332.
- [248] Levy H, Horak DA, Tegtmeier BR, et al. The value of bronchoalveolar lavage and bronchial washings in the diagnosis of invasive pulmonary aspergillosis[J]. Respir Med, 1992,86: 243-248.
- [249] Kahn FW, Jones JM, England DM. The role of bronchoalveolar lavage in the diagnosis of invasive pulmonary aspergillosis[J]. Am J Clin Pathol, 1986, 86(4): 518-523.
- [250] Guo YL, Chen YQ, Wang K, et al. Accuracy of BAL galactomannan in diagnosing invasive aspergillosis: a bivariate metaanalysis and systematic review[J]. Chest, 2010, 138: 817-824. DOI: 10.1378/chest.10-0488.
- [251] Meersseman W, Lagrou K, Maertens J, et al. Galactomannan in bronchoalveolar lavage fluid: a tool for diagnosing aspergillosis in intensive care unit patients[J]. Am J Respir Crit Care Med, 2008, 177: 27-34. DOI: 10.1164 / rccm. 200704-606OC.
- [252] Mattei D, Rapezzi D, Mordini N, et al. False-positive Aspergillus galactomannan enzyme-linked immunosorbent assay results in vivo during amoxicillin-clavulanic acid treatment[J]. J Clin Microbiol, 2004, 42: 5362-5363. DOI: 10.1128/JCM.42.11.5362-5363.2004.
- [253] Adam O, Auperin A, Wilquin F, et al. Treatment with piperacillin-tazobactam and false-positive Aspergillus galactomannan antigen test results for patients with hematological malignancies[J]. Clin Infect Dis, 2004, 38: 917-920. DOI: 10.1086/383148.
- [254] Viscoli C, Machetti M, Cappellano P, et al. False-positive galactomannan platelia Aspergillus test results for patients receiving piperacillin-tazobactam[J]. Clin Infect Dis, 2004,38: 913-916. DOI: 10.1086/382224.
- [255] Aubry A, Porcher R, Bottero J, et al. Occurrence and kinetics of false-positive Aspergillus galactomannan test results following treatment with beta-lactam antibiotics in patients with hematological disorders[J]. J Clin Microbiol, 2006, 44:

- 389-394. DOI: 10.1128/JCM.44.2.389-394.2006.
- [256] Musher B, Fredricks D, Leisenring W, et al. Aspergillus galactomannan enzyme immunoassay and quantitative PCR for diagnosis of invasive aspergillosis with bronchoalveolar lavage fluid[J]. J Clin Microbiol, 2004,42: 5517-5522. DOI: 10.1128/JCM.42.12.5517-5522.2004.
- [257] Lim WS, Baudouin SV, George RC, et al. BTS guidelines for the management of community acquired pneumonia in adults: update 2009[J]. Thorax, 2009, 64(Suppl 3): iii1-55. DOI: 10.1136/thx.2009.121434.
- [258] Mandell LA, Wunderink RG, Anzueto A, et al. Infectious Diseases Society of America / American Thoracic Society consensus guidelines on the management of community-acquired pneumonia in adults[J]. Clin Infect Dis, 2007, 44(Suppl 2): S27-72. DOI: 10.1086/511159.
- [259] Gibson SP, Weir DC, Bruge PS. A prospective audit of the value of fibre optic bronchoscopy in adults admitted with community acquired pneumonia[J]. Respir Med, 1993, 87(2): 105-109.
- [260] Manali E, Papadopoulos A, Tsiodras S, et al. The impact on community acquired pneumonia empirical therapy of diagnostic bronchoscopic techniques[J]. Scand J Infect Dis, 2008,40: 286-292. DOI: 10.1080/00365540701663373.
- [261] Jimenez P, Sldias F, Meneses M, et al. Diagnostic fiberoptic bronchoscopy in patients with community-acquired pneumonia. Comparison between bronchoalveolar lavage and telescoping plugged catheter cultures[J]. Chest, 1993, 103(4): 1023-1027.
- [262] Jimenez P, Meneses M, Saldias F, et al. Pneumococcal antigen detection in bronchoalveolar lavage fluid from patients with pneumonia[J]. Thorax, 1994,49: 872-874.
- [263] Jacobs JA, Stobberingh EE, Cornelissen EI, et al. Detection of Streptococcus pneumoniae antigen in bronchoalveolar lavage fluid samples by a rapid immunochromatographic membrane assay[J]. J Clin Microbiol, 2005, 43: 4037-4040. DOI: 10.1128/JCM.43.8.4037-4040.2005.
- [264] Jaulhac B, Nowicki M, Bornstein N, et al. Detection of Legionella spp. In bronchoalveolar lavage fluids by DNA amplification[J]. J Clin Microbiol, 1992, 30: 920-924.
- [265] Reischl U, Linde HJ, Lehn N, et al. Direct detection and differentiation of Legionella spp. And Legionella pneumophila in clinical specimens by dual-color real-time PCR and melting curve analysis[J]. J Clin Microbiol, 2002, 40: 3814-3817. DOI: 10.1128/JCM.40.10.3814-3817.2002.
- [266] Ghamande S, Rafanan A, Dweik R, et al. Role of transbronchial needle aspiration in patients receiving mechanical ventilation[J]. Chest, 2002, 122: 985-989. DOI: 10.1378/chest.122.3.985.
- [267] Perkins GD, Chatterjee S, McAuley DF, et al. Role of nonbronchoscopic lavage for investigating alveolar inflammation and permeability in acute respiratory distress syndrome[J]. Crit Care Med, 2006,34: 57-64. DOI: 10.1097/01.CCM.0000190197.69945.
- [268] Steinberg KP, Mitchell DR, Maunder RJ, et al. Safety of bronchoalveolar lavage in patients with adult respiratory distress syndrome[J]. Am Rev Respir Dis, 1993, 148: 556-561. DOI: 10.1164/ajrccm/148.3.556.
- [269] Bauer TT, Torres A, Ewig S, et al. Effects of bronchoalveolar lavage volume on arterial oxygenation in mechanically ventilated patients with pneumonia[J]. Intensive Care Med, 2001, 27: 384-393.
- [270] Hilbert G, Gruson D, Vargas F, et al. Bronchoscopy with bronchoalveolar lavage via the laryngeal mask airway in high-risk hypoxemic immunosuppressed patients[J]. Crit Care Med, 2001, 29: 249-255.
- [271] Klein U, Karzai W, Zimmermann P, et al. Changes in pulmonary mechanics after fiberoptic bronchoalveolar lavage in mechanically ventilated patients[J]. Intensive Care Med, 1998,24: 1289-1293.
- [272] Maitre B, Jaber S, Maggiore SM, et al. Continuous positive airway pressure during fiberoptic bronchoscopy in hypoxemic patients. A randomized double-blind study using a new device [J]. Am J Respir Crit Care Med, 2000,162(3 Pt 1): 1063-1067. DOI: 10.1164/ajrccm.162.3.9910117.
- [273] Papazian L, Colt HG, Scemama F, et al. Effects of consecutive protected specimen brushing and bronchoalveolar lavage on gas exchange and hemodynamics in ventilated patients[J]. Chest, 1993,104: 1548-1552. DOI:10.1378/chest.104.5.1548.
- [274] Previgliano IJ, Ripoll PI, Chiappero G, et al. Optimizing cerebral perfusion pressure during fiberoptic bronchoscopy in severe head injury: effect of hyperventilation[J]. Acta Neurochir Suppl, 2002,81: 103-105.
- [275] Trouillet JL, Guiguet M, Gibert C, et al. Fiberoptic bronchoscopy in ventilated patients. Evaluation of cardiopulmonary risk under midazolam sedation[J]. Chest, 1990,97: 927-933. DOI:10.1378/chest.97.4.927.
- [276] MacIntyre NR, Ramage JE, Follett JV. Jet Ventilation in support of fiberoptic bronchoscopy[J]. Crit Care Med, 1987,15: 303-307.
- [277] Antonelli M, Conti G, Riccioni L, et al. Noninvasive positive-pressure ventilation via face mask during bronchoscopy with ABL in high-risk hypoxemic patients[J]. Chest, 1996, 110(3): 724-728. DOI:10.1378/chest.110.3.724.
- [278] Chiner E, Sancho-Chust JN, Llombart M, et al. Fiberoptic bronchoscopy during nasal non-invasive ventilation in acute respiratory failure[J]. Respiration, 2010, 80: 321-326. DOI: 10.1159/000314074.
- [279] Baumann HJ, Klose H, Simon M, et al. Fiber optic bronchoscopy in patients with acute hypoxemic respiratory failure requiring noninvasive ventilation—a feasibility study [J]. Crit Care, 2011,15: R179. DOI: 10.1186/cc10328.
- [280] Brimacombe J, Dunbar-Reid K. The effect of introducing fiberoptic bronchoscopes on gas flow in laryngeal masks and tracheal tubes[J]. Anaesthesia, 1996, 51: 923-928. DOI: 10.1111/j.1365-2044.1996.tb14958.x.
- [281] Kim YH, Suh GY, Kim MH, et al. Safety and usefulness of bronchoscopy in ventilator-dependent patients with severe thrombocytopenia[J]. Anaesth Intensive Care, 2008, 36: 411-417.
- [282] Bajwa MK, Henein S, Kamholz SL. Fiberoptic bronchoscopy in the presence of space-occupying intracranial lesions[J]. Chest,1993,104: 101-103. DOI: 10.1378/chest.104.1.101.
- [283] Kerwin AJ, Croce MA, Timmons SD, et al. Effect of fiberoptic bronchoscopy on intracranial pressure in patients with brain injury: a prospective clinical study[J]. J Trauma, 2000, 48: 878-882.
- [284] Peerless JR, Snow N, Likavec MJ, et al. The effect of fiberoptic bronchoscopy on cerebral hemodynamics in patients with severe head injury[J]. Chest, 1995, 108: 962-965. DOI: 10.1378/chest.108.4.962.
- [285] Jose RJ, Shaefi S, Navani N. Anesthesia for bronchoscopy[J]. Curr Opin Anaesthesiol, 2014, 27(4): 453-457. DOI: 10.1097/

- ACO.0000000000000087.
- [286] Haenel JB, Moore FA, Moore EE, et al. Efficacy of selective intrabronchial air insufflation in acute lobar collapse[J]. *Am J Surg*, 1992, 164: 501-505. DOI: 10.1016 / S0002-9610(05) 81189-4.
 - [287] Lee TS, Wright BD. Selective insufflation of collapsed lung with fiberoptic bronchoscope and Swan-Ganz catheter[J]. *Intensive Care Med*, 1981,7: 241-243.
 - [288] Tsao TC, Tsai YH, Lan RS, et al. Treatment for collapsed lung in critically ill patients. Selective intrabronchial air insufflation using the fiberoptic bronchoscope[J]. *Chest*, 1990, 97: 435-438. DOI: 10.1378/chest.97.2.435.
 - [289] Tabboush ZS, Ayash RH, Badran HM. When fiberoptic bronchoscopy is indicated in the management of postoperative atelectasis[J]. *Acta Anaesthesiol Scand*, 1998, 42: 384. DOI: 10.1111/J.1399-6576.1998.tb04935.x.
 - [290] Jawarski A, Goldberg SK, Walenstein MD, et al. Utility of immediate postlobectomy fiberoptic bronchoscopy in preventing atelectasis[J]. *Chest*, 1988, 94: 38-43. DOI: 10.1378/chest.94.1.38.
 - [291] Barreiro B, Dorca J, Manresa F, et al. Protected bronchoalveolar lavage in the diagnosis of ventilator-associated pneumonia[J]. *Eur Respir J*, 1996, 9: 1500-1507. DOI:10.1183/09031936.96.09071500.
 - [292] Bello S, Tajada A, Chacon E, et al. 'Blind' protected specimen brushing versus bronchoscopic techniques in the aetiological diagnosis of ventilator-associated pneumonia[J]. *Eur Respir J*, 1996, 9: 1494-1499. DOI: 10.1183 / 09031936.96.09071494.
 - [293] Casetta M, Bolt F, Antoun S, et al. Diagnosis of nosocomial pneumonia in cancer patients undergoing mechanical ventilation: a prospective comparison of the plugged telescoping catheter with the protected specimen brush[J]. *Chest*, 1999, 115: 1641-1645. DOI: 10.1378/chest.115.6.1641.
 - [294] Humphreys H, Winter R, Baker M, et al. Comparison of bronchoalveolar lavage and catheter lavage to confirm ventilator-associated lower respiratory tract infection[J]. *J Med Microbiol*, 1996, 45: 226-231. DOI: 10.1099/00222615- 45-3-226.
 - [295] Jorda R, Parras F, Ibanez J, et al. Diagnosis of nosocomial pneumonia in mechanically ventilated patients by the blind protected telescoping catheter[J]. *Intensive Care Med*, 1993, 19: 377-382.
 - [296] Jourdain B, Novara A, Joly-Guillou ML, et al. Role of quantitative cultures of endotracheal aspirates in the diagnosis of nosocomial pneumonia[J]. *Am J Respir Crit Care Med*, 1995, 152: 241-246. DOI: 10.1164/ajrcm.152.1.7599831.
 - [297] Kirtland SH, Corley DE, Winterbauer RH, et al. The diagnosis of ventilator-associated pneumonia: a comparison of histologic, microbiologic, and clinical criteria[J]. *Chest*, 1997, 112: 445-457. DOI: 10.1378/chest.112.2.445.
 - [298] Leal-Noval SR, Alfora-Rodriguez E, Murillo-Cabeza F, et al. Diagnostic value of the blind brush in mechanically ventilated patients with nosocomial pneumonia[J]. *Intensive Care Med*, 1992, 18: 410-414.
 - [299] Marquette CH, Herengt F, Mathieu D, et al. Diagnosis of pneumonia in mechanically ventilated patients. Repeatability of the protected specimen brush[J]. *Am Rev Respir Dis*, 1993, 147: 211-214. DOI: 10.1164/ajrcm/147.1.211.
 - [300] Marquette CH, Herengt F, Saulnier F, et al. Protected specimen brush in the assessment of ventilator-associated pneumonia. Selection of a certain lung segment for bronchoscopic sampling is unnecessary[J]. *Chest*, 1993, 103: 243-247. DOI: 10.1378/chest.103.1.243.
 - [301] Pugin J, Auckenthaler R, Mili N, et al. Diagnosis of ventilator-associated pneumonia by bacteriologic analysis of bronchoscopic and nonbronchoscopic 'blind' bronchoalveolar lavage fluid[J]. *Am Rev Respir Dis*, 1991, 143(5 Pt 1): 1121-1129. DOI: 10.1164/ajrcm/143.5_Pt_1.1121.
 - [302] Rouby JJ, Rossignon MD, Nicolas MH, et al. A prospective study of protected bronchoalveolar lavage in the diagnosis of nosocomial pneumonia[J]. *Anesthesiology*, 1989, 71: 679-685.
 - [303] Sanchez-Nieto JM, Torres A, Garcia-Cordoba F, et al. Impact of invasive and noninvasive quantitative culture sampling on outcome of ventilator-associated pneumonia: a pilot study[J]. *Am J Respir Crit Care Med*, 1998, 157: 371-376. DOI: 10.1164/ajrcm.157.2.97-02039.
 - [304] Souweine B, Veber B, Bedos JP, et al. Diagnostic accuracy of protected specimen brush and bronchoalveolar lavage in nosocomial pneumonia: impact of previous antimicrobial treatments[J]. *Crit Care Med*, 1998, 26: 236-244.
 - [305] Swanson JM, Wood GC, Croce MA, et al. Utility of preliminary bronchoalveolar lavage results in suspected ventilator-associated pneumonia[J]. *J Trauma*, 2008, 65: 1271-1277. DOI: 10.1097/TA.0b013e3181574d6a.
 - [306] Berton DC, Kalil AC, Teiveira PJ. Quantitative versus qualitative cultures of respiratory secretions for clinical outcomes in patients with ventilator-associated pneumonia[J]. *Cochrane Database Syst Rev*, 2012, (1): CD006482. DOI: 10.1002/14651858.CD006482.pub3.
 - [307] Collins SR, Blank RS. Fiberoptic intubation: an overview and update[J]. *Respir Care*, 2014, 59(6): 865-878; discussion 878-880. DOI: 10.4187/respcare.03012.
 - [308] Rodrigues AJ, Scordamaglio PR, Palomino AM, et al. Difficult airway intubation with flexible bronchoscope[J]. *Braz J Anesthesiol*, 2013, 63(4): 358-361. DOI: 10.1016 / j. bjane.2012.05.001.
 - [309] Gill N, Purohit S, Kalra P, et al. Comparison of hemodynamic responses to intubation: Flexible fiberoptic bronchoscope versus McCoy laryngoscope in presence of rigid cervical collar simulating cervical immobilization for traumatic cervical spine [J]. *Anesth Essays Res*, 2015, 9(3): 337-342. DOI: 10.4103/ 0259-1162.158013.
 - [310] Asai M, Samayoa AX, Hodge C, et al. Elective intubation and positive pressure ventilation for transbronchial lung biopsy[J]. *J Surg Res*, 2017, 219: 296-301. DOI: 10.1016 / j. jss.2017.05.085.
 - [311] Bulpa PA, Dive AM, Mertens L, et al. Combined bronchoalveolar lavage and transbronchial lung biopsy: safety and yield in ventilated patients[J]. *Eur Respir J*, 2003, 21(3): 489-494. DOI:10.1183/09031936.03.00298303
 - [312] 农凌波, 李时悦, 何为群, 等. 机械通气下经支气管镜肺活检[J]. *中国呼吸与危重症监护杂志*, 2009, 8(3): 275-278. DOI: 10.3969/j.issn.1671-6205.2009.03.017.
 - [313] Bermasconi M, Chhajed PN, Muller P, et al. Patients' satisfaction with flexible bronchoscopy in a hospital-based community practice[J]. *Respiration*, 2009, 78: 440-445. DOI: 10.1159/000228906.
 - [314] Department of Health UK. Understanding what matters: a guide to using patient feedback to transform care. May 2009. http://webarchive.nationalarchives.gov.uk/+/www.dh.gov.uk/en/Publicationsandstatistics/Publications/PublicationsPolicyAndGuidance/DH_099780.

- [315] D'Ippolito R, Foresi A, Castagnetti C, et al. Indications for flexible fiberoptic bronchoscopy and its safety in the very elderly[J]. Monaldi Arch ChestDis, 2007, 67: 23-29. DOI: 10.4081/monaldi.2007.506.
- [316] 张杰, 董淑文. 支气管镜操作中镇静镇痛药物应用的评价[J]. 中华结核和呼吸杂志, 2010, 33(9): 709-711. DOI: 10.3760/cma.j.issn.1001-0939.2010.09.023.
- [317] Crawford M, Pollock J, Anderson K, et al. Comparison of midazolam with propofol for sedation in outpatient bronchoscopy[J]. Br J Anaesth, 1993, 70(4): 419-422.
- [318] 刘云喜, 邢玉斌, 巩玉秀, 等. 软式内镜清洗消毒技术规范[D]. 中华人民共和国国家卫生和计划生育委员会. <http://www.nhpc.gov.cn/zhuz/s9496/201701/491ec38efc884531801549cfb90d865d.shtml>.
- [319] Beilenhoff U, Neumann CS, Rey JF, et al. The ESGE Guidelines Committee. ESGE-ESCEA guideline: cleaning and disinfection in gastrointestinal endoscopy[J]. Update 2008. Endoscopy, 2008, 40: 939-957. DOI: 10.1055/s-2008-1077722.
- [320] National Endoscopy Programme. Decontamination standards for flexible endoscopes 2008. 2009. [http://www.thejag.org.uk/downloads%5CUit%20Resources%5CDecontamination%20Standards%20for%10Flexible%20Endoscopy.pdf](http://www.thejag.org.uk/downloads%5CUnit%20Resources%5CDecontamination%20Standards%20for%10Flexible%20Endoscopy.pdf) (accessed Feb 2019).
- [321] MHRA. Medical Devices Agency (MDA) Device Bulletin DB2002 (05) on decontamination of endoscopes. 2005. <http://www.mhra.gov.uk/Publications/Safetyguidance/DeviceBulletins/CON007329>.
- [322] The Health Act 2006: Code of practice for the prevention and control of healthcare associated infections. 2006. <http://www.hpa.org.uk/Topics/InfectiousDiseases/InfectionsAZ/HCAI/GuidelinesForProfessionalsHCAI/> (accessed Feb 2019).
- [323] Health and Safety Executive (HSE) UK. An evaluation of disinfecting agents used in endoscopy suites. 2007. <http://www.hse.gov.uk/research/rrhtm/rr445.htm> (accessed Feb 2019).
- [324] British Society of Gastroenterology Provision of Endoscopy Related Services in District General Hospitals. 2001. http://www.bsg.org.uk/images/stories/docs/clinical/guidelines/endoscopy/endo_related_services.pdf (accessed Feb 2019).
- [325] 嵇凤官. 《电离辐射防护与辐射源安全基本标准》GB18871-2002 介绍[J]. 核标准计量与质量, 2004, (4): 41-48.
- [326] 潘自强(校译). 国际放射防护委员会(ICRP)2007 年 103 号文件[M]. 原子能出版社, 2008.
- [327] CIRSE, Cardiovascular and Interventional Radiological Society of Europe. Occupational radiation protection in interventional radiology: a joint guideline of the Cardiovascular and Interventional Radiology Society of Europe and the Society of Interventional Radiology[J]. J Vasc Interv Radiol, 2010, 21: 607-615. DOI:10.1016/j.jvir.2010.01.007.

(收稿日期: 2019-04-01)

(本文编辑: 吕小东)

·读者·作者·编者·

本刊“介入园地”栏目征稿

近年来,随着介入呼吸病学的迅速发展,国内外针对呼吸内镜技术的相关研究不断拓展和深入,已成为呼吸病学中一个朝气蓬勃、前景广阔的新领域。借助荧光支气管镜、超声引导下的经支气管镜淋巴结活检、电烧灼、冷冻、气道内支架置入、球囊扩张和光动力治疗等呼吸内镜相关新兴技术,呼吸系统疾病的诊断和治疗手段有了长足进步,但目前我国介入呼吸病学的发展水平与发达国家相比还有一定差距,介入技术的普及程度仍然不足,更重要的是介入技术的应用尚缺乏规范。

为宣传、普及、探讨和逐步规范介入呼吸病学技术的应用,提供一个供相关专业人员交流、争鸣以及相互学习的平台,本刊自 2010 年起开辟“介入园地”栏目,来稿形式不拘,以临床报道为主,无论是学科最新进展,还是疑难和(或)经典病例介绍,或是临床经验总结及临床实践中所遇到的问题,均欢迎踊跃赐稿。

来稿需通过中华医学会远程稿件处理系统(www.cma.org.cn)上传,作者投稿操作说明可通过中华医学会网站业务中心下载。

投稿时请务必注明作者的联系电话(手机)、单位(包括科室)及电子邮箱,稿件格式参见本刊已刊登的文章格式。